CHALMERS
FINITE ELEMENT CENTER

PREPRINT 2001–15

A Finite Element Method for Domain Decomposition with Non-Matching Grids

Roland Becker, Peter Hansbo and Rolf Stenberg



Chalmers Finite Element Center
CHALMERS UNIVERSITY OF TECHNOLOGY
Göteborg Sweden 2001

CHALMERS FINITE ELEMENT CENTER

Preprint 2001–15

A Finite Element Method for Domain Decomposition with Non-Matching Grids

Roland Becker, Peter Hansbo and Rolf Stenberg

CHALMERS

Chalmers Finite Element Center
Chalmers University of Technology
SE–412 96 Göteborg Sweden
Göteborg, August 2001

A Finite Element Method for Domain Decomposition with Non-Matching Grids
Roland Becker, Peter Hansbo and Rolf Stenberg
NO 2001–15
ISSN 1404–4382

Chalmers Finite Element Center
Chalmers University of Technology
SE–412 96 Göteborg
Sweden
Telephone: +46 (0)31 772 1000
Fax: +46 (0)31 772 3595
www.phi.chalmers.se

Printed in Sweden
Chalmers University of Technology
Göteborg, Sweden 2001

A FINITE ELEMENT METHOD FOR DOMAIN DECOMPOSITION WITH NON-MATCHING GRIDS

ROLAND BECKER, PETER HANSBO, AND ROLF STENBERG

Abstract. In this note, we propose and analyse a method for handling interfaces between non-matching grids based on an approach suggested by Nitsche [15] for the approximation of Dirichlet boundary conditions. The exposition is limited to self-adjoint elliptic problems, using Poisson's equation as a model. A priori and a posteriori error estimates are given. Some numerical results are included.

1. Introduction

In any domain decomposition method, one has to define how the continuity between the subdomains is to be enforced. Different approaches have been proposed:

- Iterative procedures, enforcing that the approximate solution or its normal derivative or combinations thereof should be continuous across interfaces. This forms the basis for the standard Schwarz alternating method as defined, e.g., by Lions [14].
- Direct procedures, using Lagrange multiplier techniques to achieve continuity. Different variants have been proposed, e.g., by Le Tallec and Sassi [13], and Bernadi, Maday, and Patera [8].

The multiplier method has the advantage of directly yielding a solvable global system. However, in the latter method, new unknowns (the multipliers) must be introduced and solved for. The method must then either satisfy the inf-sup condition, which necessitates special choices of multiplier spaces (such as mortar elements, cf. [8]), or then stabilization techniques (cf. Baiocchi, Brezzi, and Marini [3]) must be used.

In this paper, we consider a third possibility, i.e. Nitsche's method [15], which was originally introduced for the purpose of solving Dirichlet problems without enforcing the boundary conditions in the definition of the finite element spaces. This method has later been used by Arnold [2] for the discretization of second order elliptic equations by discontinuous finite elements. In earlier papers [16, 17] we have pointed out the close connection between Nitsche's method and stabilized methods and proposed it as a mortaring method.

Date: August 30, 2001.

Key words and phrases. Nitsche's method, domain decomposition, non-matching grids.

Roland Becker, Institute of Applied Mathematics, Heidelberg, INF 294, D-69120 Heidelberg, Germany, *email*: roland.becker@iwr.uni-heidelberg.de

Peter Hansbo, Department of Solid Mechanics, Chalmers University of Technology, S–412 96 Göteborg, Sweden, *email*: hansbo@math.chalmers.se

Rolf Stenberg, Institute of Mathematics, Box 1100, FIN-02015 Helsinki University of Technology,Finland *email*: rolf.stenberg@hut.fi.

In this paper we will give a more detailed analysis of this domain decomposition technique where independent approximations are used on the different subdomains. The continuity of the solution across interfaces is enforced weakly, but in such a way that the resulting discrete scheme is consistent with the original partial differential equation. Under some regularity assumptions we derive both a priori and a posteriori error estimates. We also give numerical results obtained with the method.

Although we discuss its application to domain decomposition, the same technique is also suited for other applications, e.g.,

- to handle diffusion terms in the discontinuous Galerkin method [2, 6];
- to simplify mesh generation (different parts can be meshed independently from each other);
- finite element methods with different polynomial degree on adjacent elements;
- new finite element methods such as linear approximations on quadrilaterals.

2. The domain decomposition method

In this section we will introduce the mortaring method based on the classical method of Nitsche. We will perform a classical stability and a priori error analysis. For simplicity, we consider the model Poisson problem, i.e. of solving the partial differential equation

$$\begin{aligned} -\Delta u &= f \quad \text{in } \Omega, \\ u &= 0 \quad \text{on } \partial\Omega, \end{aligned} \tag{2.1}$$

where Ω is a bounded domain in two or three space dimensions and $f \in L_2(\Omega)$.

Likewise for ease of presentation, we consider only the case where Ω is divided into two non-overlapping subdomains Ω_1 and Ω_2, $\Omega_1 \cup \Omega_2$, with interface $\Gamma = \overline{\Omega}_1 \cap \overline{\Omega}_2$. We further assume that the subdomains are polyhedral (or polygonal in $\mathbb{R}^2$) and that Γ is polygonal (or a broken line).

This equation can be written in weak form as: find $u \in H_0^1(\Omega)$ such that

$$a(u,v) = (f,v), \quad \forall v \in H_0^1(\Omega), \tag{2.2}$$

where

$$a(u,v) = \int_\Omega \nabla u \cdot \nabla v \, dx, \quad (f,v) = \int_\Omega f\, v \, dx, \tag{2.3}$$

and $H_0^1(\Omega)$ is the space of square-integrable functions, with square-integrable first derivatives, that vanish on the boundary $\partial\Omega$ of Ω.

Our discrete method for the approximate solution of (2.1) is a nonconforming finite element method which is continuous within each Ω_i and discontinuous across Γ. We start

by rewriting the original problem (2.1) as two equations and the interface conditions:

$$\begin{aligned} -\Delta u_i &= f \quad \text{in } \Omega_i, \quad i=1,2, \\ u_i &= 0 \quad \text{on } \partial\Omega \cap \Omega_i, \quad i=1,2, \\ u_1 - u_2 &= 0 \quad \text{on } \Gamma, \\ \frac{\partial u_1}{\partial n_1} + \frac{\partial u_2}{\partial n_2} &= 0 \quad \text{on } \Gamma. \end{aligned} \tag{2.4}$$

Here, n_i is the outward unit normal to $\partial\Omega_i$.

We will perform our analysis under the following regularity assumption.

Assumption 2.1. *The solution of* (2.2) *satisfies* $u \in H^s(\Omega)$*, with* $s > 3/2$.

With this assumption it holds $\partial u_i/\partial n_i \in L_2(\Gamma)$ and the two problems (2.1) and (2.4) are equivalent (see for example [1]) with:

$$u|_{\Omega_i} = u_i, \quad i=1,2. \tag{2.5}$$

In the following we will therefore write $u = (u_1, u_2) \in V_1 \times V_2$ with the continuous spaces

$$V_i = \left\{ v_i \in H^1(\Omega_i) : \ \partial v_i/\partial n_i \in L_2(\Gamma), \ v_i|_{\partial\Omega\cap\partial\Omega_i} = 0 \right\}, \quad i=1,2.$$

To formulate our method, we suppose that we have regular finite element partitionings $\mathbf{T}_h^i$ of the subdomains Ω_i into shape regular simplexes. These two meshes induce "trace" meshes on the interface

$$\mathbf{G}_h^i = \{ \ E \ : E = K \cap \Gamma, \ K \in \mathbf{T}_h^i \ \}. \tag{2.6}$$

By h_K and h_E we denote the diameter of element $K \in \mathbf{T}_h^i$ and $E \in \mathbf{G}_h^i$, respectively. For the purpose of the a priori analysis, we also define

$$h = \max\{h_K, \ h_E \ : \ K \in \mathbf{T}_h^i, \ E \in \mathbf{G}_h^i, \ i=1,2. \ \}.$$

We seek the approximation $u_h = (u_{1,h}, u_{2,h})$ in the space $V^h = V_1^h \times V_2^h$, where

$$V_i^h = \left\{ v_i \in V_i : \ v_i|_K \text{ is a polynomial of degree } p \text{ for all } K \in \mathbf{T}_h^i \right\},$$

choosing for simplicity p constant on all subdomains. On the interface we will use the notation

$$[\![v]\!] := v_1 - v_2 \tag{2.7}$$

for the jump,

$$\{ v \} := \frac{1}{2} v_1 + \frac{1}{2} v_2 \tag{2.8}$$

for the average and

$$\langle v, w \rangle_S := \int_S v \, w \, ds, \qquad \text{for } S \in \mathbf{G}_h^i, \text{ or } S = \Gamma. \tag{2.9}$$

Finally, we denote

$$n := n_1 = -n_2. \tag{2.10}$$

With this notation we have

$$\left\{\frac{\partial v}{\partial n}\right\} = \frac{1}{2}\frac{\partial v_1}{\partial n_1} - \frac{1}{2}\frac{\partial v_2}{\partial n_2} \tag{2.11}$$

and hence

$$\left\{\frac{\partial u}{\partial n}\right\} = \frac{\partial u}{\partial n} = \frac{\partial u_1}{\partial n_1} = -\frac{\partial u_2}{\partial n_2}. \tag{2.12}$$

The methods of Nitsche [15] and Arnold [2] now give the following domain decomposition method.

Mortar Method. *Find $u_h \in V^h$ such that*

$$a_h(u_h, v) = f_h(v) \quad \forall v \in V^h, \tag{2.13}$$

with

$$\begin{aligned} a_h(w,v) \;:=\; & \sum_{i=1}^{2}\left[(\nabla w_i, \nabla v_i)_{\Omega_i} + \gamma \sum_{E\in\mathbf{G}_h^i} h_E^{-1}\left\langle [\![w]\!], [\![v]\!]\right\rangle_E\right] \\ & - \left\langle \left\{\frac{\partial w}{\partial n}\right\}, [\![v]\!]\right\rangle_\Gamma - \left\langle \left\{\frac{\partial v}{\partial n}\right\}, [\![w]\!]\right\rangle_\Gamma \end{aligned} \tag{2.14}$$

and

$$f_h(v) := \sum_{i=1}^{2}(f, v_i)_{\Omega_i}, \tag{2.15}$$

where $\gamma > 0$ is chosen sufficiently large, see Lemma 2.5 below.

The first observation is that this formulation gives a consistent method.

Lemma 2.2. *The solution $u = (u_1, u_2)$ to (2.4) satisfies*

$$a_h(u, v) = f_h(v) \quad \forall v \in V. \tag{2.16}$$

Proof. Multiplying the first equation in (2.4) with v_i, integrating over Ω_i, using Greens formula and the relations (2.12) yields

$$\begin{aligned} f_h(v) \;&=\; \sum_{i=1}^{2}(f, v_i)_{\Omega_i} = \sum_{i=1}^{2}\left[(\nabla u_i, \nabla v_i)_{\Omega_i} - \left\langle \frac{\partial u_i}{\partial n_i}, v_i\right\rangle_\Gamma\right] \\ &=\; \sum_{i=1}^{2}(\nabla u_i, \nabla v_i)_{\Omega_i} - \left\langle \left\{\frac{\partial u}{\partial n}\right\}, [\![v]\!]\right\rangle_\Gamma. \end{aligned} \tag{2.17}$$

Since $[\![u]\!] = 0$ on Γ we have

$$0 = -\left\langle \left\{\frac{\partial v}{\partial n}\right\}, [\![u]\!]\right\rangle_\Gamma + \gamma \sum_{i=1}^{2}\sum_{E\in\mathbf{G}_h^i} h_E^{-1}\left\langle [\![u]\!], [\![v]\!]\right\rangle_E. \tag{2.18}$$

Adding (2.17) and (2.18) the gives the claim. ∎

For the stability analysis below we need the following mesh-dependent dual norms. To emphasize that Γ is to be considered as a part of $\partial\Omega_i$ we write Γ_i.

$$\|v\|^2_{1/2,h,\Gamma_i} = \sum_{E\in\mathbf{G}^i_h} h_E^{-1}\ \|v\|^2_{L_2(E)} \tag{2.19}$$

and

$$\|v\|^2_{-1/2,h,\Gamma_i} = \sum_{E\in\mathbf{G}^i_h} h_E\ \|v\|^2_{L_2(E)}\,, \tag{2.20}$$

which satisfy

$$|\,\langle v,w\rangle_\Gamma\,| \le \|v\|_{1/2,h,\Gamma_i}\ \|v\|_{-1/2,h,\Gamma_i}\,. \tag{2.21}$$

We further define the norms

$$\|v\|^2_{1,h,\Omega_i} = \|\nabla v_i\|^2_{L_2(\Omega_i)} + \|[\![v]\!]\|^2_{1/2,h,\Gamma_i}\,, \tag{2.22}$$

$$\|v\|^2_{1,h} = \sum_{i=1}^{2} \|v_i\|^2_{1,h,\Omega_i}, \tag{2.23}$$

and

$$|||v|||^2_{1,h,\Omega_i} = \|v\|^2_{1,h,\Omega_i} + \left\|\frac{\partial v_i}{\partial n_i}\right\|^2_{-1/2,h,\Gamma_i}\,, \tag{2.24}$$

$$|||v|||^2_{1,h} = \sum_{i=1}^{2} |||v_i|||^2_{1,h,\Omega_i}. \tag{2.25}$$

The following estimate is readily proved by local scaling.

Lemma 2.3. *There is a positive constant C_I such that*

$$\left\|\frac{\partial v_i}{\partial n_i}\right\|^2_{-1/2,h,\Gamma_i} \le C_I\,\|\nabla v_i\|^2_{L^2(\Omega_i)} \quad \forall v_i\in V_i^h. \tag{2.26}$$

For linear elements ∇v is constant on each element and then it is particularly easy to give a bound for the constant C_I, see Remark 2.11 below.

A immediate consequence of the above lemma is the equivalence of norms in the finite element subspace.

Lemma 2.4. *The norms $\|\cdot\|_{1,h}$ and $|||\cdot|||_{1,h}$ are equivalent on the subspace V^h.*

The stability of the method can now be proved.

Lemma 2.5. *Suppose that $\gamma > C_I/4$. Then it holds*

$$a_h(v,v) \ge C|||v|||^2_{1,h} \qquad \forall v\in V^h.$$

Proof. From the definition (2.14), the relation (2.12), and the preceeding lemmas we get

$$\begin{aligned}
a_h(v,v) &= \sum_{i=1}^{2} \Big[\|\nabla v_i\|^2_{L_2(\Omega_i)} + \gamma \|[\![v]\!]\|^2_{1/2,h,\Gamma_i} \Big] - 2 \left\langle [\![v]\!], \left\{ \frac{\partial v}{\partial n} \right\} \right\rangle_\Gamma \\
&= \sum_{i=1}^{2} \Big[\|\nabla v_i\|^2_{L_2(\Omega_i)} + \gamma \|[\![v]\!]\|^2_{1/2,h,\Gamma_i} \Big] - \left\langle [\![v]\!], \frac{\partial v_1}{\partial n_1} \right\rangle_\Gamma + \left\langle [\![v]\!], \frac{\partial v_2}{\partial n_2} \right\rangle_\Gamma \\
&\geq \sum_{i=1}^{2} \left[\|\nabla v_i\|^2_{L_2(\Omega_i)} + \gamma \|[\![v]\!]\|^2_{1/2,h,\Gamma_i} - \left| \left\langle [\![v]\!], \frac{\partial v_i}{\partial n_i} \right\rangle_\Gamma \right| \right] \\
&\geq \sum_{i=1}^{2} \left[\|\nabla v_i\|^2_{L_2(\Omega_i)} + \gamma \|[\![v]\!]\|^2_{1/2,h,\Gamma_i} - \|v\|_{1/2,h,\Gamma_i} \left\| \frac{\partial v_i}{\partial n_i} \right\|_{-1/2,h,\Gamma_i} \right] \\
&\geq \sum_{i=1}^{2} \Big[\big(1 - \frac{C_I}{2\varepsilon}\big) \|\nabla v_i\|^2_{L_2(\Omega_i)} + (\gamma - \frac{\varepsilon}{2}) \|[\![v]\!]\|^2_{1/2,h,\Gamma_i} \Big] \\
&\geq C_1 \|v\|^2_{1,h} \\
&\geq C_2 |||v|||^2_{1,h},
\end{aligned}$$

for $\gamma > \varepsilon/2$ and choosing $\varepsilon > C_I/2$. ∎

The following interpolation estimates holds, cf. Thomée [18].

Lemma 2.6. *Suppose that $u \in H^s(\Omega)$, with $3/2 < s \leq p+1$. Then it holds*

$$\inf_{v \in V^h} \|u - v\|_{1,h} \leq C h^{s-1} \|u\|_{H^s(\Omega)} \tag{2.27}$$

and

$$\inf_{v \in V^h} |||u - v|||_{1,h} \leq C h^{s-1} \|u\|_{H^s(\Omega)}. \tag{2.28}$$

We are now able to prove the following a priori error estimate.

Theorem 2.7. *Suppose $\gamma > C_I/4$. Then it holds*

$$|||u - u_h|||_{1,h} \leq C \inf_{v \in V^h} |||u - v|||_{1,h}.$$

If $u \in H^s(\Omega)$, for $3/2 < s \leq p+1$, then

$$|||u - u_h|||_{1,h} \leq C h^{s-1} \|u\|_{H^s(\Omega)}.$$

Proof. Since the bilinear form a_h is bounded with respect to the norm $|||\cdot|||_{1,h}$, the first inequality follows from the stability estimate of Lemma 2.5 and the triangle inequality. The second estimate then follows from Lemma 2.6. ∎

Remark 2.8. *As $[\![u]\!] = 0$ on Γ the error estimate gives $\| [\![u_h]\!] \|_{-1/2,h,\Gamma_i} = O(h^{s-1})$. For quasi-uniform meshes it then holds $\| [\![u_h]\!] \|_{L_2(\Gamma)} = O(h^{s-1/2})$.*

Remark 2.9. *The presented method resembles a mesh-dependent penalty method, but with added consistency terms involving normal derivatives across the interface. Note that the formulation allows us to deduce optimal order error estimates with preserved condition number of $O(h^{-2})$ for a quasiuniform mesh. Pure penalty methods, in contrast, are not consistent, and optimal error estimates require degrading the condition number for higher polynomial approximation (cf.* [5]*).*

Remark 2.10. *The form $a_h(\cdot,\cdot)$ in (2.14) is symmetric and positive definite, which is natural as the problem to be approximated has the same properties. With this there exists fast solvers for the resulting matrix problem. If the bilinear form is changed to*

$$
\begin{aligned}
a_h(w,v) \;:=\; & \sum_{i=1}^{2}\Big[(\nabla w_i,\nabla v_i)_{\Omega_i} + \gamma \sum_{E\in\mathbf{G}_h^i} h_E^{-1}\left\langle [\![w]\!],[\![v]\!]\right\rangle_E\Big] \\
& - \left\langle \left\{\frac{\partial w}{\partial n}\right\},[\![v]\!]\right\rangle_\Gamma + \left\langle \left\{\frac{\partial v}{\partial n}\right\},[\![w]\!]\right\rangle_\Gamma ,
\end{aligned}
\tag{2.29}
$$

then one obtains a method which is stable for all positive values of γ:

$$
a_h(v,v) \geq C|||v|||_{1,h}^2 \qquad \forall v\in V^h,\ \forall\gamma>0.
$$

If the Laplace operator is a part of a problem which is not symmetric, then it might be practical to use this nonsymmetric bilinear form. See [10, 11] *for applications to convection diffusion problems.*

Remark 2.11. *For linear elements ∇v_i ($v_i\in V_i^h$) is constant on each element and hence for $K\in\mathbf{T}_h^i$ with the base $E\in\mathbf{G}_h^i$ we have*

$$
\left\|\frac{\partial v_i}{\partial n_i}\right\|_{0,E}^2 = meas(E)\left|\frac{\partial v_i}{\partial n_i}\right|^2 \tag{2.30}
$$

and

$$
\|\nabla v_i\|_{0,K}^2 \geq \left\|\frac{\partial v_i}{\partial n_i}\right\|_{0,K}^2 = meas(K)\left|\frac{\partial v_i}{\partial n_i}\right|^2 \tag{2.31}
$$

and hence it holds

$$
h_E\left\|\frac{\partial v_i}{\partial n_i}\right\|_{0,E}^2 \leq \frac{h_E\, meas(E)}{meas(K)}\,\|\nabla v_i\|_{0,K}^2 . \tag{2.32}
$$

Hence, once the shape regularity of the mesh is specified, one has a bound for C_I. In a practical implementation easier would be to replace the weight γh_E^{-1} in (2.14) with a parameter

$$
\alpha_K = \alpha\,\frac{meas(E)}{meas(K)}, \tag{2.33}
$$

with $\alpha>1/4$ fixed. With this all results of the paper hold.

3. A posteriori error estimates

We will first consider control of the error $e = u - u_h$ in the mesh dependent energy norm $|||\cdot|||_{1,h}$. We define the local and global estimators as

$$E_K(u_h)^2 = h_K^2 \|f + \Delta u_h\|_{L_2(K)}^2 + h_K \left\| \left[\left[\frac{\partial u_h}{\partial n_K} \right]\right] \right\|_{L_2(\partial K)}^2 + h_K^{-1} \|[\![u_h]\!]\|_{L_2(\partial K)}^2 \tag{3.1}$$

and

$$E(u_h) = \left[\sum_{K \in \mathbf{T}_h} E_K(u_h)^2 \right]^{1/2}. \tag{3.2}$$

To be able to control the normal derivatives across the interface, we introduce a "saturation" assumption similar to the one used by Wohlmuth [19] in the context of a posteriori error estimates for the mortar element method. (We also remark that the assumption is consistent with the interpolation estimates of Theorem 2.6).

Assumption 3.1. *There is a constant C such that*

$$\left\| \frac{\partial (u_i - u_{i,h})}{\partial n_i} \right\|_{-1/2,h,\Gamma_i} \leq C \|u_i - u_{i,h}\|_{1,h,\Omega_i}. \tag{3.3}$$

A consequence of this assumption is the bound

$$|||u - u_h|||_{1,h} \leq C \|u - u_h\|_{1,h}. \tag{3.4}$$

The a posteriori estimate is now the following.

Theorem 3.2. *Suppose that the Assumption 3.1 is valid. Then there is a positive constant C such that*

$$|||u - u_h|||_{1,h} \leq C E(u_h). \tag{3.5}$$

Proof. We denote $e = u - u_h$. By Lemma 2.2 we have

$$\begin{aligned} \|e\|_{1,h}^2 &= a_h(e,e) + 2 \left\langle [\![e]\!], \left\{ \frac{\partial e}{\partial n} \right\} \right\rangle_\Gamma + (1-\gamma) \sum_{i=1}^{2} \sum_{E \in \mathbf{G}_h^i} h_E^{-1} \langle [\![e]\!], [\![e]\!] \rangle_E \\ &:= R_1 + R_2, \end{aligned} \tag{3.6}$$

with

$$R_1 := a_h(e,e),$$

and

$$R_2 := 2 \left\langle [\![e]\!], \left\{ \frac{\partial e}{\partial n} \right\} \right\rangle_\Gamma + (1-\gamma) \sum_{i=1}^{2} \|[\![e]\!]\|_{1/2,h,\Gamma_i}^2.$$

Since $[\![e]\!]|_\Gamma = [\![u_h]\!]|_\Gamma$ Assumption 3.1 gives

$$\begin{aligned} R_2 &= \left\langle [\![u_h]\!], \frac{\partial e_1}{\partial n_1} \right\rangle_\Gamma - \left\langle [\![u_h]\!], \frac{\partial e_2}{\partial n_2} \right\rangle_\Gamma + (1-\gamma) \sum_{i=1}^{2} \|[\![e]\!]\|^2_{1/2,h,\Gamma_i} \\ &\leq C \sum_{i=1}^{2} (\|[\![u_h]\!]\|_{1/2,h,\Gamma_i} \left\| \frac{\partial e_i}{\partial n_i} \right\|_{-1/2,h,\Gamma_i} + \|[\![u_h]\!]\|_{1/2,h,\Gamma_i} \|[\![e]\!]\|_{1/2,h,\Gamma_i}) \\ &\leq C\|e\|_{1,h} \Big(\sum_{i=1}^{2} \|[\![u_h]\!]\|_{1/2,h,\Gamma_i} \Big) \leq C\|e\|_{1,h} E(u_h). \end{aligned} \tag{3.7}$$

Next, let $\pi_h e$ be the Clément interpolant to e, which satisfies

$$\Big(\sum_{K \in \mathbf{T}_h} \Big(h_K^{-2} \|e - \pi_h e\|^2_{L_2(K)} + h_K^{-1} \|e - \pi_h e\|^2_{L_2(\partial K)} \Big) \Big)^{1/2} \leq C\|e\|_{1,h} \tag{3.8}$$

and

$$\|e - \pi_h e\|_{1,h} \leq C\|e\|_{1,h}. \tag{3.9}$$

From the consistency (2.2) we have $a_h(e, \pi_h e) = 0$. Hence

$$\begin{aligned} R_1 &= a_h(e,e) = a_h(e, e - \pi_h e) \\ &= \sum_{i=1}^{2} \Big((\nabla e_i, \nabla(e_i - \pi_h e_i))_{\Omega_i} + \gamma \sum_{E \in \mathbf{G}_h^i} h_E^{-1} \langle [\![e]\!], [\![e - \pi_h e]\!] \rangle_E \Big) \\ &\quad - \left\langle \left\{ \frac{\partial e}{\partial n} \right\}, [\![e - \pi_h e]\!] \right\rangle_\Gamma - \left\langle \left\{ \frac{\partial (e - \pi_h e)}{\partial n} \right\}, [\![e]\!] \right\rangle_\Gamma \\ &:= S_1 + S_2 + S_3 + S_4. \end{aligned} \tag{3.10}$$

Integrating by parts on each $K \in \mathbf{T}_h$ yields

$$\begin{aligned} S_1 &= \sum_{i=1}^{2} (\nabla e_i, \nabla(e_i - \pi_h e_i))_{\Omega_i} \\ &= \sum_{K \in \mathbf{T}_h} \Big(-(\Delta e, e - \pi_h e)_K + \langle \frac{\partial e}{\partial n_K}, e - \pi_h e \rangle_{\partial K} \Big) \\ &:= T_1 + T_2. \end{aligned} \tag{3.11}$$

Since on $K \in \mathbf{T}_h$ it holds $-\Delta e = -\Delta u + \Delta u_h = f + \Delta u_h$, the first term above is estimated using (3.8)

$$
\begin{aligned}
T_1 &= -\sum_{K\in\mathbf{T}_h} (\Delta e, e-\pi_h e)_K \\
&\leq \Big(\sum_{K\in\mathbf{T}_h} h_K^2 \left\|\Delta u_h + f\right\|_{L_2(K)}^2\Big)^{1/2}\Big(\sum_{K\in\mathbf{T}_h} h_K^{-2} \left\|e-\pi_h e\right\|_{L_2(K)}^2\Big)^{1/2} \\
&\leq \Big(\sum_{K\in\mathbf{T}_h} h_K^2 \left\|\Delta u_h + f\right\|_{L_2(K)}^2\Big)^{1/2} \|e\|_{1,h}.
\end{aligned} \tag{3.12}
$$

Let $\mathbf{I}_h$ be the collection of element sides in the interiors of the subdomains Ω_i. The boundary integrals in (3.11) above we now split into those in $\mathbf{I}_h$ and those lying on the interface Γ:

$$
\begin{aligned}
T_2 = \sum_{K\in\mathbf{T}_h} \langle \frac{\partial e}{\partial n_K}, e-\pi_h e\rangle_{\partial K} = \\
\sum_{E\in\mathbf{I}_h} \langle \frac{\partial e}{\partial n_E}, e-\pi_h e\rangle_E + \langle \frac{\partial e_1}{\partial n_1}, e_1-\pi_h e_1\rangle_\Gamma + \langle \frac{\partial e_2}{\partial n_2}, e_2-\pi_h e_2\rangle_\Gamma.
\end{aligned} \tag{3.13}
$$

The integrals over the interior sides can now be grouped together two by two yielding the estimate

$$
\begin{aligned}
&|\sum_{E\in\mathbf{I}_h} \langle \frac{\partial e}{\partial n_E}, e-\pi_h e\rangle_E| \\
&\leq C\Big[\sum_{K\in\mathbf{T}_h} h_K \Big\|\Big[\Big[\frac{\partial u_h}{\partial n_K}\Big]\Big]\Big\|_{L_2(\partial K)}^2\Big]^{1/2}\Big[\sum_{K\in\mathbf{T}_h} h_K^{-1}\left\|e-\pi_h e\right\|_{L_2(\partial K)}^2\Big]^{1/2} \\
&\leq C\Big[\sum_{K\in\mathbf{T}_h} h_K \Big\|\Big[\Big[\frac{\partial u_h}{\partial n_K}\Big]\Big]\Big\|_{L_2(\partial K)}^2\Big]^{1/2}\|e\|_{1,h},
\end{aligned} \tag{3.14}
$$

where (3.8) was used.

Using the definition (2.11) we have

$$
\begin{aligned}
S_3 &= -\Big\langle \Big\{\frac{\partial e}{\partial n}\Big\}, [\![e-\pi_h e]\!]\Big\rangle_\Gamma = -\Big\langle \frac{1}{2}\frac{\partial e_1}{\partial n_1} + \frac{1}{2}\frac{\partial e_2}{\partial n_2}, e_1-\pi_h e_1 - (e_2-\pi_h e_2)\Big\rangle_\Gamma \\
&= -\Big\langle \frac{1}{2}\frac{\partial e_1}{\partial n_1} - \frac{1}{2}\frac{\partial e_2}{\partial n_2}, e_1-\pi_h e_1\Big\rangle_\Gamma + \Big\langle \frac{1}{2}\frac{\partial e_1}{\partial n_1} - \frac{1}{2}\frac{\partial e_2}{\partial n_2}, (e_2-\pi_h e_2)\Big\rangle_\Gamma .
\end{aligned} \tag{3.15}
$$

Combining these terms with two last terms in (3.13) gives

$$
\begin{aligned}
&\langle\frac{\partial e_1}{\partial n_1}, e_1-\pi_h e_1\rangle_\Gamma + \langle\frac{\partial e_2}{\partial n_2}, e_2-\pi_h e_2\rangle_\Gamma - \left\langle\left\{\frac{\partial e}{\partial n}\right\}, [\![e-\pi_h e]\!]\right\rangle_\Gamma \\
&= \left\langle(1-\frac{1}{2})\frac{\partial e_1}{\partial n_1}+\frac{1}{2}\frac{\partial e_2}{\partial n_2}, e_1-\pi_h e_1\right\rangle_\Gamma + \left\langle\frac{1}{2}\frac{\partial e_1}{\partial n_1}+(1-\frac{1}{2})\frac{\partial e_2}{\partial n_2}, (e_2-\pi_h e_2)\right\rangle_\Gamma \\
&= \frac{1}{2}\left\langle\frac{\partial e_1}{\partial n_1}+\frac{\partial e_2}{\partial n_2}, e_1-\pi_h e_1\right\rangle_\Gamma + \frac{1}{2}\left\langle\frac{\partial e_1}{\partial n_1}+\frac{\partial e_2}{\partial n_2}, e_2-\pi_h e_2\right\rangle_\Gamma \\
&= \frac{1}{2}\left\langle\frac{\partial (e_1-e_2)}{\partial n}, e_1-\pi_h e_1\right\rangle_\Gamma + \frac{1}{2}\left\langle\frac{\partial (e_1-e_2)}{\partial n}, e_2-\pi_h e_2\right\rangle_\Gamma \\
&= \frac{1}{2}\left\langle\left[\!\left[\frac{\partial e}{\partial n}\right]\!\right], e_1-\pi_h e_1\right\rangle_\Gamma + \frac{1}{2}\left\langle\left[\!\left[\frac{\partial e}{\partial n}\right]\!\right], e_2-\pi_h e_2\right\rangle_\Gamma \qquad (3.16)\\
&= \frac{1}{2}\left\langle\left[\!\left[\frac{\partial u_h}{\partial n}\right]\!\right], e_1-\pi_h e_1\right\rangle_\Gamma + \frac{1}{2}\left\langle\left[\!\left[\frac{\partial u_h}{\partial n}\right]\!\right], e_2-\pi_h e_2\right\rangle_\Gamma \\
&\le \frac{1}{2}E(u_h)\Big[\sum_{K\in\mathbf{T}_h} h_K^{-1}\,\|e-\pi_h e\|^2_{L_2(\partial K)}\Big]^{1/2} \\
&\le CE(u_h)\|e\|_{1,h}.
\end{aligned}
$$

From (3.11) to (3.16) we now get

$$S_1 + S_3 \le C\|e\|_{1,h}E(u_h). \tag{3.17}$$

From the Clément estimate (3.8) have

$$S_2 = \gamma\sum_{i=1}^{2}\sum_{E\in\mathbf{G}_h^i} h_E^{-1}\,\langle[\![e]\!], [\![e-\pi_h e]\!]\rangle_E \le C\|e\|_{1,h}E(u_h).$$

To estimate the last term S_4 we use Lemma 2.3 and the Assumption 3.1

$$
\begin{aligned}
S_4 &= -\left\langle\left\{\frac{\partial(e-\pi_h e)}{\partial n}\right\}, [\![e]\!]\right\rangle_\Gamma = -\left\langle\left\{\frac{\partial e}{\partial n}\right\}, [\![e]\!]\right\rangle_\Gamma + \left\langle\left\{\frac{\partial \pi_h e}{\partial n}\right\}, [\![e]\!]\right\rangle_\Gamma \\
&\le C\|e\|_{1,h}E(u_h). \qquad (3.18)
\end{aligned}
$$

Hence, collecting the estimates (3.10) to (3.18) yields

$$R_1 \le C\|e\|_{1,h}E(u_h), \tag{3.19}$$

which together with (3.7) proves the estimate

$$\|u-u_h\|_{1,h} = \|e\|_{1,h} \le CE(u_h). \tag{3.20}$$

The claim then follows from (3.4). ∎

Next, we consider the error in the L_2-norm. This is measured with the estimator

$$L(u_h) = \Big[\sum_{K\in\mathbf{T}_h} h_K^2 E_K(u_h)^2\Big]^{1/2}. \tag{3.21}$$

For the a posteriori estimate we as usual, need the H^2-regularity but not Assumption 3.1.

Theorem 3.3. *Let z be the solution to the problem*

$$\begin{aligned} -\Delta z &= g \quad \text{in } \Omega, \\ z &= 0 \quad \text{on } \partial\Omega, \end{aligned} \tag{3.22}$$

and suppose that the shift theorem

$$\|z\|_{H^2(\Omega)} \leq C \|g\|_{L_2(\Omega)} \tag{3.23}$$

is valid. Then there is a positive constant C such that

$$\|u - u_h\|_{L_2(\Omega)} \leq CL(u_h). \tag{3.24}$$

Proof. We again write $z = (z_1, z_2)$ and note that

$$\frac{\partial z_1}{\partial n_1} + \frac{\partial z_2}{\partial n_2} = 0 \quad \text{and} \quad \left\{ \frac{\partial z}{\partial n} \right\} = \frac{\partial z_1}{\partial n_1} = -\frac{\partial z_2}{\partial n_2}.$$

Choosing $g = u - u_h$ in (3.22) we then get

$$\begin{aligned} \|u - u_h\|^2_{L_2(\Omega)} &= \sum_{i=1}^{2} \Big[(\nabla z_i, \nabla(u_i - u_{i,h}))_{\Omega_i} - \langle \frac{\partial z_i}{\partial n_i}, u_i - u_{i,h} \rangle_{\Gamma_i} \Big] \\ &= \sum_{i=1}^{2} \Big[(\nabla z_i, \nabla(u_i - u_{i,h}))_{\Omega_i} + \langle \frac{\partial z_i}{\partial n_i}, u_{i,h} \rangle_{\Gamma_i} \Big] \\ &= \sum_{i=1}^{2} (\nabla z_i, \nabla(u_i - u_{i,h}))_{\Omega_i} + \langle \left\{ \frac{\partial z}{\partial n} \right\}, [\![u_h]\!] \rangle_{\Gamma}. \end{aligned} \tag{3.25}$$

We let $\pi_h z \in V^h$ be the Clément interpolant to z satisfying

$$\begin{aligned} &\Big(\sum_{K \in \mathbf{T}_h} \Big(h_K^{-4} \|z - \pi_h z\|^2_{L_2(K)} + h_K^{-3} \|z - \pi_h z\|^2_{L_2(\partial K)} + h_K^{-1} \left\| \frac{\partial (z - \pi_h z)}{\partial n_K} \right\|^2_{L_2(\partial K)} \Big) \Big)^{1/2} \\ &\leq C \|z\|_{H^2(\Omega)}. \end{aligned} \tag{3.26}$$

From the consistency condition we have $a_h(u-u_h,\pi_h z)=0$, and since $[\![z]\!]=0,\ [\![u]\!]=0$, on the interface, we get

$$
\begin{aligned}
0 &= a_h(u-u_h,\pi_h z) \\
&= \sum_{i=1}^{2}\Big[(\nabla(u_i-u_{i,h}),\nabla\pi_h z_i)_{\Omega_i}+\gamma\sum_{E\in\mathbf{G}_h^i}h_E^{-1}\left\langle[\![u-u_h]\!],[\![\pi_h z]\!]\right\rangle_E\Big] \\
&\quad -\left\langle\left\{\frac{\partial(u-u_h)}{\partial n}\right\},[\![\pi_h z]\!]\right\rangle_\Gamma-\left\langle\left\{\frac{\partial\pi_h z}{\partial n}\right\},[\![u-u_h]\!]\right\rangle_\Gamma \\
&= \sum_{i=1}^{2}\Big[(\nabla(u_i-u_{i,h}),\nabla\pi_h z_i)_{\Omega_i}+\gamma\sum_{E\in\mathbf{G}_h^i}h_E^{-1}\left\langle[\![\ u_h]\!],[\![z-\pi_h z]\!]\right\rangle_E\Big] \\
&\quad +\left\langle\left\{\frac{\partial(u-u_h)}{\partial n}\right\},[\![z-\pi_h z]\!]\right\rangle_\Gamma+\left\langle\left\{\frac{\partial\pi_h z}{\partial n}\right\},[\![u_h]\!]\right\rangle_\Gamma.
\end{aligned}
\tag{3.27}
$$

Substracting (3.27) from (3.25) gives

$$
\begin{aligned}
&\|u-u_h\|_{L_2(\Omega)}^2 \\
&=\sum_{i=1}^{2}\Big[(\nabla(z_i-\pi_h z,\nabla(u_i-u_{i,h}))_{\Omega_i}-\gamma\sum_{E\in\mathbf{G}_h^i}h_E^{-1}\left\langle[\![\ u_h]\!],[\![z-\pi_h z]\!]\right\rangle_E\Big] \\
&\quad -\left\langle\left\{\frac{\partial(u-u_h)}{\partial n}\right\},[\![z-\pi_h z]\!]\right\rangle_\Gamma+\left\langle\left\{\frac{\partial(z-\pi_h z)}{\partial n}\right\},[\![u_h]\!]\right\rangle_\Gamma \\
&:= R_1+R_2+R_3+R_4.
\end{aligned}
\tag{3.28}
$$

Using Schwarz inequality and the Clément estimate (3.26) we get

$$
\begin{aligned}
&R_2+R_4 \\
&\le C\Big(\sum_{K\in\mathbf{T}_h}\Big(h_K^{-3}\|z-\pi_h z\|_{L_2(\partial K)}^2+h_K^{-1}\left\|\frac{\partial(z-\pi_h z)}{\partial n_K}\right\|_{L_2(\partial K)}^2\Big)\Big)^{1/2}\Big(\sum_{E\in\mathbf{G}_h^i}h_E\|[\![u_h]\!]\|_{L_2(E)}\Big)^{1/2} \\
&\le C\|z\|_{H^2(\Omega)}L(u_h)\le C\|u-u_h\|_{L_2(\Omega)}L(u_h).
\end{aligned}
$$

Integrating by parts yields

$$
\begin{aligned}
R_1 + R_3 &= \sum_{i=1}^{2} \Big[(\nabla(z_i - \pi_h z, \nabla(u_i - u_{i,h}))_{\Omega_i} - \left\langle \left\{ \frac{\partial(u - u_h)}{\partial n} \right\}, [\![z - \pi_h z]\!] \right\rangle_{\Gamma} \\
&= \sum_{K \in \mathbf{T}_h} \left\{ (z - \pi_h z, \Delta(u_h - u))_K + \left\langle \frac{\partial(u - u_h)}{\partial n_K}, z - \pi_h z \right\rangle_{\partial K} \right\} - \left\langle \left\{ \frac{\partial(u - u_h)}{\partial n} \right\}, [\![z - \pi_h z]\!] \right\rangle_{\Gamma} \\
&= \sum_{K \in \mathbf{T}_h} (z - \pi_h z, \Delta u_h + f)_K + \sum_{E \in \mathbf{I}_h} \left\langle \frac{\partial(u - u_h)}{\partial n_E}, z - \pi_h z \right\rangle_E \\
&\quad + \sum_{i=1}^{2} \left\langle \frac{\partial(u_i - u_{i,h})}{\partial n_i}, z_i - \pi_h z_i \right\rangle_{\Gamma} - \left\langle \left\{ \frac{\partial(u - u_h)}{\partial n} \right\}, [\![z - \pi_h z]\!] \right\rangle_{\Gamma}.
\end{aligned}
$$

The boundary terms in the second term above are grouped together two by two yielding jump terms in the normal derivative. This gives the estimate

$$
\begin{aligned}
&\sum_{K \in \mathbf{T}_h} (z - \pi_h z, \Delta u_h + f)_K + \sum_{E \in \mathbf{I}_h} \left\langle \frac{\partial(u - u_h)}{\partial n_E}, z - \pi_h z \right\rangle_E \\
&\leq C \Bigg(\sum_{K \in \mathbf{T}_h} \left(h_K^{-4} \left\| z - \pi_h z \right\|_{L_2(K)}^2 + h_K^{-3} \left\| z - \pi_h z \right\|_{L_2(\partial K)}^2 \right) \Bigg)^{1/2} \\
&\qquad \times \Bigg(\sum_{K \in \mathbf{T}_h} h_K^4 \left\| f + \Delta u_h \right\|_{L_2(K)}^2 + h_K^3 \left\| \left[\!\!\left[\frac{\partial u_h}{\partial n_K} \right]\!\!\right] \right\|_{L_2(\partial K)}^2 \Bigg)^{1/2} \\
&\leq C \left\| z \right\|_{H^2(\Omega)} L(u_h) \leq C \left\| u - u_h \right\|_{L_2(\Omega)} L(u_h).
\end{aligned} \tag{3.29}
$$

The remaining two terms are estimated as follows

$$
\begin{aligned}
&\sum_{i=1}^{2}\left\langle \frac{\partial(u_i-u_{i,h})}{\partial n_i}, z_i-\pi_h z_i\right\rangle_\Gamma - \left\langle \left\{\frac{\partial(u-u_h)}{\partial n}\right\}, [\![z-\pi_h z]\!]\right\rangle_\Gamma \\
&= \sum_{i=1}^{2}\left\langle \frac{\partial u_i}{\partial n_i}, z_i-\pi_h z_i\right\rangle_\Gamma - \sum_{i=1}^{2}\left\langle \frac{\partial u_{i,h}}{\partial n_i}, z_i-\pi_h z_i\right\rangle_\Gamma \\
&\qquad - \left\langle \frac{\partial u}{\partial n}, [\![z-\pi_h z]\!]\right\rangle_\Gamma + \left\langle \left\{\frac{\partial u_h}{\partial n}\right\}, [\![z-\pi_h z]\!]\right\rangle_\Gamma \\
&= -\sum_{i=1}^{2}\left\langle \frac{\partial u_{i,h}}{\partial n_i}, z_i-\pi_h z_i\right\rangle_\Gamma + \left\langle \left\{\frac{\partial u_h}{\partial n}\right\}, [\![z-\pi_h z]\!]\right\rangle_\Gamma \\
&= -\left\langle \frac{\partial u_{1,h}}{\partial n_1}, z_1-\pi_h z_1\right\rangle_\Gamma - \left\langle \frac{\partial u_{2,h}}{\partial n_2}, z_2-\pi_h z_2\right\rangle_\Gamma \\
&\qquad +\frac{1}{2}\left\langle \frac{\partial u_{1,h}}{\partial n_1}, z_1-\pi_h z_1\right\rangle_\Gamma - \frac{1}{2}\langle \frac{\partial u_{1,h}}{\partial n_1}, z_2-\pi_h z_2\right\rangle_\Gamma \\
&\qquad -\frac{1}{2}\left\langle \frac{\partial u_{2,h}}{\partial n_2}, z_1-\pi_h z_1\right\rangle_\Gamma + \frac{1}{2}\left\langle \frac{\partial u_{2,h}}{\partial n_2}, z_2-\pi_h z_2\right\rangle_\Gamma \\
&= -\frac{1}{2}\left\langle \frac{\partial u_{1,h}}{\partial n_1}, z_1-\pi_h z_1\right\rangle_\Gamma - \frac{1}{2}\left\langle \frac{\partial u_{2,h}}{\partial n_2}, z_2-\pi_h z_2\right\rangle_\Gamma \\
&\qquad -\frac{1}{2}\left\langle \frac{\partial u_{1,h}}{\partial n_1}, z_2-\pi_h z_2\right\rangle_\Gamma - \frac{1}{2}\langle \frac{\partial u_{2,h}}{\partial n_2}, z_1-\pi_h z_1\right\rangle_\Gamma \\
&= -\frac{1}{2}\left\langle \frac{\partial u_{1,h}}{\partial n_1}+\frac{\partial u_{2,h}}{\partial n_2}, z_1-\pi_h z_1\right\rangle_\Gamma - \frac{1}{2}\left\langle \frac{\partial u_{1,h}}{\partial n_1}+\frac{\partial u_{2,h}}{\partial n_2}, z_2-\pi_h z_2\right\rangle_\Gamma \\
&= -\frac{1}{2}\left\langle \left[\!\left[\frac{\partial u_h}{\partial n}\right]\!\right], z_1-\pi_h z_1\right\rangle_\Gamma - \frac{1}{2}\left\langle \left[\!\left[\frac{\partial u_h}{\partial n}\right]\!\right], z_2-\pi_h z_2\right\rangle_\Gamma \\
&\le CL(u_h)\Big(\sum_{K\in\mathbf{T}_h} h_K^{-3}\left\|z-\pi_h z\right\|^2_{L_2(\partial K)}\Big)^{1/2} \\
&\le CL(u_h)\left\|z\right\|_{H^2(\Omega)} \le CL(u_h)\left\|u-u_h\right\|_{L_2(\Omega)}.
\end{aligned}
\tag{3.30}
$$

Combining (3.29) and (3.30) gives

$$R_1 + R_3 \le CL(u_h)\left\|u-u_h\right\|_{L_2(\Omega)}$$

which together with (3.29) and (3.28) proves the claim. ∎

4. Numerical examples

4.1. **Numerical verification of the a priori estimates.** To verify the a priori estimates, we choose the model problem of a unit square with exact solution

$$u = x\,y\,(1-x)\,(1-y)$$

corresponding to a right-hand side of $f = 2(x - x^2 + y - y^2)$. The domain is divided by a vertical slit at $x = 0.7$. Two different triangulations were used: one matching and one non-matching, see Figure 1.

In Figure 2 (left-hand side) we give the convergence in the broken energy norm. The dashed line is the non-matching grid computation. Both meshes show the same convergence with slope 0.95. which is close to the theoretical value of 1. On the right-hand side we show the convergence of the L_2-norm of the jump term (dashed line for the non-matching grid). Here we obtain a better convergence (slope 2.15) for the matching grids than for the non-matching grids (slope 1.57, close to the theoretical value of 3/2).

4.2. **Adaptive computations.** We present results of adaptive computations on the L-shaped domain

$$\Omega = (0,1) \times (0,1) \setminus (1/2,1) \times (0,1/2).$$

The problem is boundary driven ($f = 0$), with boundary data corresponding to the exact solution $u = r^{2/3} \sin(2\theta/3)$ in polar coordinates (with origin at $(1/2, 1/2)$). We let $\Omega_1 = (0, 1/2) \times (0, 1)$ and $\Omega_2 = (1/2, 1) \times (1/2, 1)$, and use a non-matching triangulation. The purpose of this example is not to obtain exact error control, but rather to show how the adaptive algorithm behaves with respect to the elements adjacent to the interface. We consider adaptive control of the L_2-norm error, but we have not made any attempt to measure the constant in the inequality (3.23); instead we have simply tuned the interpolation constants to approximately match the exact error.

In Figure 3 and 4 we show the first and last (adapted) meshes resulting from equilibrating the error distribution over the set of elements (for details, see [12, 7]). In Figure 5 we show the exact and estimated L_2-errors on the sequence of meshes, which show a reasonable agreement. For more exact error control, more computational effort must be invested.

4.3. **Difference between matching and non-matching meshes.** In the previous example, the adaptive algorithm produced a slightly finer mesh on the interface. One natural question is then whether the effect of non-matching actually does lead to larger errors. In Figure 6 and 7 we show two different meshes for solving the problem described in Section 4.1, and the corresponding nodal interpolants of the errors. The interface is sitauted at $x = 1/2$, and it is noted that the error on the interface is markedly larger than in the interior of the domains only in the case of non-matching meshes.

References

[1] J.-P. Aubin, *Approximation of Elliptic Boundary-Value Problem.* Wiley (1972).

[2] D. Arnold, An interior penalty finite element method with discontinuous elements. *SIAM J. Num. Anal.* **19** (1982) 742–760.

[3] C. Baiocchi, F. Brezzi, and L. D. Marini, Stabilization of Galerkin methods and applications to domain decomposition. A. Bensoussan and J.-P. Verjus Eds., *Future Tendencies in Computer Science, Control and Applied Mathematics,* Springer (1992) 345–355.

[4] J. C. Barbosa and T. J. R. Hughes, Boundary Lagrange multipliers in finite element methods: error analysis in natural norms. *Numer. Math.* **62** (1992) 1–15.

[5] J. W. Barrett and C. M. Elliot, Finite element approximation of the Dirichlet problem using the boundary penalty method. *Numer. Math.* **49** (1986) 343–366.

[6] R. Becker and P. Hansbo, Discontinuous Galerkin methods for convection-diffusion problems with arbitrary Péclet number. P. Neittaanmäki, T. Tiihonen, and P. Tarvainen Eds., *Numerical Mathematics and Advanced Applications: Proceedings of the 3rd European Conference*, World Scientific (2000) 100-109.

[7] R. Becker and R. Rannacher, A feed-back approach to error control in finite element methods: basic analysis and examples. *East-West J. Numer. Math.* **4** (1996) 237–264.

[8] C. Bernadi, Y. Maday, and A. Patera, A new nonconforming approach to domain decomposition: the mortar element method, in: *Nonlinear Partial Differential Equations and Their Application,* H. Brezis and J. L. Lions (eds.), (Pitman, 1989).

[9] F. Brezzi, L.P. Franca, D. Marini, and A. Russo, Stabilization techniques for domain decomposition methods with non-matching grids, IAN-CNR Report N. 1037, Istituto di Analisi Numerica Pavia.

[10] J. Freund, R. Stenberg, On weakly imposed boundary conditions for second order problems. M. Morandi Cecchi et al. Eds., *Proceedings of the Ninth Int. Conf. Finite Elements in Fluids,* (Venice 1995) 327-336

[11] J. Freund, Space-time finite element methods for second order problems: an algorithmic approach. *Acta Polytech. Scand. Mathe. Comput. Manage. Engrg. Ser.* 79 (1996).

[12] C. Johnson and P. Hansbo, Adaptive finite element methods in computational mechanics. *Comput. Methods Appl. Mech. Engrg.* **101** (1992) 143–181.

[13] P. Le Tallec and T. Sassi, Domain decomposition with nonmatching grids: augmented Lagrangian approach. *Math. Comp.* **64** (1995) 1367–1396.

[14] P. L. Lions, On the Schwarz alternating method III: a variant for nonoverlapping subdomains. T. F. Chan, R. Glowinski, J. Periaux, and O. B. Widlund Eds., *Third International Symposium on Domain Decomposition Methods for Partial Differential Equations,* SIAM (1989) 202–223.

[15] J. Nitsche, Über ein Variationsprinzip zur Lösung von Dirichlet-Problemen bei Verwendung von Teilräumen, die keinen Randbedingungen unterworfen sind. *Abh. Math. Univ. Hamburg* **36** (1970) 9–15.

[16] R. Stenberg, On some techniques for approximating boundary conditions in the finite element method. *J. Comput. Appl. Math.* **63** (1995) 139–148.

[17] R. Stenberg, Mortaring by a method of J.A. Nitsche. S Idelsohn, E. Onate and E. Dvorkin Eds., *Computational Mechanics: New Trends and Applications,* CIMNE, Barcelona (1998).

[18] V. Thomée, *Galerkin Finite Element Methods for Parabolic Problems*, (Springer, 1997)

[19] B. I. Wohlmuth, A residual based error estimator for mortar finite element discretizations. *Numer. Math.* **84** (1999) 143–171.



FIGURE 1. Matching and non-matching grids.



FIGURE 2. Convergence in energy and convergence of the jump term.



FIGURE 3. First mesh and final adapted mesh.



FIGURE 4. Elevation of the solution.



FIGURE 5. Estimated and exact errors in the $L_2(\Omega)-$norm.



FIGURE 6. Matching meshes and nodal errors.



FIGURE 7. Non-matching meshes and nodal errors.

Chalmers Finite Element Center Preprints

2000–01 *Adaptive Finite Element Methods for the Unsteady Maxwell's Equations*
Johan Hoffman

2000–02 *A Multi-Adaptive ODE-Solver*
Anders Logg

2000–03 *Multi-Adaptive Error Control for ODEs*
Anders Logg

2000–04 *Dynamic Computational Subgrid Modeling* (Licentiate Thesis)
Johan Hoffman

2000–05 *Least-Squares Finite Element Methods for Electromagnetic Applications* (Licentiate Thesis)
Rickard Bergström

2000–06 *Discontinuous Galerkin Methods for Incompressible and Nearly Incompressible Elasticity by Nitsche's Method*
Peter Hansbo and Mats G. Larson

2000–07 *A Discountinuous Galerkin Method for the Plate Equation*
Peter Hansbo and Mats G. Larson

2000–08 *Conservation Properties for the Continuous and Discontinuous Galerkin Methods*
Mats G. Larson and A. Jonas Niklasson

2000–09 *Discontinuous Galerkin and the Crouzeix-Raviart element: Application to elasticity*
Peter Hansbo and Mats G. Larson

2000–10 *Pointwise A Posteriori Error Analysis for an Adaptive Penalty Finite Element Method for the Obstacle Problem*
Donald A. French, Stig Larson and Ricardo H. Nochetto

2000–11 *Global and Localised A Posteriori Error Analysis in the Maximum Norm for Finite Element Approximations of a Convection-Diffusion Problem*
Mats Boman

2000–12 *A Posteriori Error Analysis in the Maximum Norm for a Penalty Finite Element Method for the Time-Dependent Obstacle Problem*
Mats Boman

2000–13 *A Posteriori Error Analysis in the Maximum Norm for Finite Element Approximations of a Time-Dependent Convection-Diffusion Problem*
Mats Boman

2001–01 *A Simple Nonconforming Bilinear Element for the Elasticity Problem*
Peter Hansbo and Mats G. Larson

2001–02 *The $\mathcal{LL}^*$ Finite Element Method and Multigrid for the Magnetostatic Problem*
Rickard Bergström, Mats G. Larson, and Klas Samuelsson

2001–03 *The Fokker-Planck Operator as an Asymptotic Limit in Anisotropic Media*
Mohammad Asadzadeh

2001–04 *A Posteriori Error Estimation of Functionals in Elliptic Problems: Experiments*
Mats G. Larson and A. Jonas Niklasson

2001–05 *A Note on Energy Conservation for Hamiltonian Systems Using Continuous Time Finite Elements*
Peter Hansbo

2001–06 *Stationary Level Set Method for Modelling Sharp Interfaces in Groundwater Flow*
Nahidh Sharif and Nils-Erik Wiberg

2001–07 *Integration methods for the calculation of the magnetostatic field due to coils*
Marzia Fontana

2001–08 *Adaptive finite element computation of 3D magnetostatic problems in potential formulation*
Marzia Fontana

2001–09 *Multi-Adaptive Galerkin Methods for ODEs I: Theory & Algorithms*
Anders Logg

2001–10 *Multi-Adaptive Galerkin Methods for ODEs II: Applications*
Anders Logg

2001–11 *Energy norm a posteriori error estimation for discontinuous Galerkin methods*
Roland Becker, Peter Hansbo, and Mats G. Larson

2001–12 *Analysis of a family of discontinuous Galerkin methods for elliptic problems: the one dimensional case*
Mats G. Larson and A. Jonas Niklasson

2001–13 *Analysis of a nonsymmetric discontinuous Galerkin method for elliptic problems: stability and energy error estimates*
Mats G. Larson and A. Jonas Niklasson

2001–14 *A hybrid method for the wave equation*
Larisa Beilina, Klas Samuelsson, Krister Åhlander

2001–15 *A Finite Element Method for Domain Decomposition with Non-Matching Grids*
Roland Becker, Peter Hansbo and Rolf Stenberg

These preprints can be obtained from

`www.phi.chalmers.se/preprints`