CHALMERS
FINITE ELEMENT CENTER

PREPRINT 2004–17

Ultraconvergence of an interpolated finite element method for some fourth-order elliptic problems

Andrey B. Andreev and Milena R. Racheva



Chalmers Finite Element Center
CHALMERS UNIVERSITY OF TECHNOLOGY
Göteborg Sweden 2004

CHALMERS FINITE ELEMENT CENTER

Preprint 2004–17

Ultraconvergence of an interpolated finite element method for some fourth-order elliptic problems

Andrey B. Andreev and Milena R. Racheva

CHALMERS

Chalmers Finite Element Center
Chalmers University of Technology
SE–412 96 Göteborg Sweden
Göteborg, July 2004

Ultraconvergence of an interpolated finite element method for some fourth-order elliptic problems
Andrey B. Andreev and Milena R. Racheva
NO 2004–17
ISSN 1404–4382

Chalmers Finite Element Center
Chalmers University of Technology
SE–412 96 Göteborg
Sweden
Telephone: +46 (0)31 772 1000
Fax: +46 (0)31 772 3595
www.phi.chalmers.se

Printed in Sweden
Chalmers University of Technology
Göteborg, Sweden 2004

ULTRACONVERGENCE OF AN INTERPOLATED FINITE ELEMENT METHOD FOR SOME FOURTH-ORDER ELLIPTIC PROBLEMS

ANDREY B. ANDREEV AND MILENA R. RACHEVA

Abstract. We introduce nonstandard interpolated finite elements providing better accuracy for a fourth-order elliptic boundary value problems, as well as to the biharmonic eigenvalue problems. The term "ultraconvergence" indicates that the convergence rate is at least two orders higher than the optimal global rate. This method is a variant of a postprocessing procedure when the known finite element solution is used. Moreover, *a posteriori* error estimates of global ultraconvergent type are derived.

The presented approach is applicable for the general rectangular finite element meshes. Some numerical results illustrate the efficiency of the proposed algorithm.

1. Introduction

In recent years, superconvergence and ultraconvergence for finite element (FE) solutions have been an active area in numerical analysis and its applications. The main objective of this field of studies is to improve the existing approximation accuracy by applying certain postprocessing techniques which are relatively easy to implement.

Zienkiewicz and J. Zhu [16, 17] developed an effective error estimation technique of finite element approximations. Their method falls into the category of *recovery-based methods*. This *a posteriori* error estimation technique approximates the error in energy or an energy norm on a particular partition. There are many ideas which are closely related to the recovery technique (see [7, 13]). Mathematical proofs for the derivative patch recovery can be found in Li and Zhang [6]. For instance, there are proofs for some finite element methods under special assumptions of the partition and, in Q. Zhu and Zhao [15], for one-dimensional correction to the finite element solution. Relative references on *a posteriori* error estimation and adaptivity could be found in Ainsworth and Oden [1], and in Verfürth [11].

Our aim of this paper is to establish an ultraconvergence result for planar FE approximations of the elliptic fourth-order problem by using an interpolated FEM approach. Similar considerations are presented by Q. Lin et al. [7] for second order elliptic problems (see also [14]).

Supercloseness of a nonstandard nodal interpolant is exploited in this paper. This is done in order to develop postprocessing schemes to improve the approximation order of

1991 *Mathematics Subject Classification.* 65N30, 65N25, 65N12, 65N15, 65D10.

Key words and phrases. finite element method, ultraconvergence, superconvergence, fourth-order elliptic problems.

This work was partially supported by Technical University of Gabrovo under Contract NoIII.2/2004.

Δu_h from $\mathcal{O}(h^3)$ to $\mathcal{O}(h^5)$, especially when polynomials of degree four in each variable are applied in FEM. An ultraconvergence result for fourth-order eigenvalue problems is obtained by means of a theorem proving the supercloseness between the Ritz projection and the corresponding approximate eigenfunction.

By applying the method ultraconvergent results are obtained without using higher elements. This procedure is based on a higher interpolation of the FE solution on the original finite elements.

Here follows the general layout of the paper: In Section 2, we present a review of a model for a fourth-order problem and its finite element procedure. In Section 3, we construct an auxiliary interpolation operator i_h, which verifies certain "vertices-edges" conditions. The main body of the paper contains the proofs of superclosed properties, generated by the operator i_h. This is presented in Section 4. Section 5 is devoted to the construction of high interpolation I_{2h} and the proof of a global ultraconvergence error estimate. In Section 6, we show that the method yields an ultraconvergent scheme for a fourth-order eigenvalue problem. The paper closes by some numeric examples illustrating the theory and concluding remarks.

2. Setting of the Problem and FEM

Let us consider the fourth-order elliptic model problem:

$$\Delta^2 u(x,y) = f(x,y),\ (x,y) \in \Omega, \tag{1}$$

with boundary conditions

$$u(x,y) = \frac{\partial u}{\partial \nu}(x,y) = 0,\ (x,y) \in \Gamma, \tag{2}$$

where $\Omega \subset \mathbf{R^2}$ is a simply connected polyrectangular domain with boundary Γ and ν is the outher normal vector.

Let $H^m(\Omega)$ be the real Sobolev space for a nonnegative integer m and their associated norms $\|\cdot\|_{m,\Omega}$ and seminorms $|\cdot|_{m,\Omega}$. The weak solution of the problem (1)- (2) is a function $u(x,y)$, belonging to $V \equiv H_0^2(\Omega)$,

$$H_0^2(\Omega) = \left\{ v \in H^2(\Omega) :\ v|_\Gamma = \frac{\partial v}{\partial \nu}|_\Gamma = 0 \right\}$$

and satisfying

$$a(u,v) = (f,v)\ \forall v \in V, \tag{3}$$

where

$$a(u,v) = \int_\Omega \Delta u \Delta v\, dx\, dy$$

and the notation $(\cdot,\cdot)$ is adopted for the $L_2(\Omega)-$inner product.

Obviously the $a-$form is symmetric and coercive on V, i.e.

$$a(u,v) = a(v,u),\ \rho\|v\|_{2,\Omega}^2 \le a(v,v),\ \forall u,v \in V,$$

where ρ is a positive constant. Moreover, it is continuous:

$$|a(u,v)| \leq C\|u\|_{2,\Omega}\|v\|_{2,\Omega},\ \forall u, v \in V.$$

Henceforth, C represents generic and positive constant.

Consider a family of regular partitions τ_h of $\overline{\Omega}$ consisting of rectangular finite elements T, which fulfill standard assumptions [5] and suppose that this family satisfies the inverse assumptions ([5], p. 140). Here h is a FE parameter. We denote by Q_k the set of polynomials of degree k at most in each variable.

Let $V_h \subset V$ be finite element spaces associated with the partition τ_h. Assume that the polynomial space in the construction of V_h contains $Q_n,\ n \geq 3$. Our aim for constructing an ultraconvergent interpolant requires $n \geq 4$. In order to avoid technical difficulties we confine the considerations to the case $n = 4$.

Then

$$V_h = \left\{ v \in C^1(\overline{\Omega}) \cap V :\ v_{|T} \in Q_4(T), \forall T \in \tau_h \right\}.$$

Let $P_h :\ V \to V_h$ be the elliptic projection operator defined by

$$a(u - P_h u, v) = 0\ \forall u \in V,\ \forall v \in V_h.$$

It is well-known (see [8]) that if the solution $u(x,y)$ of (3) belongs to $H^5(\Omega) \cap V$, then

$$\|u - P_h u\|_{2,\Omega} \leq Ch^3\|u\|_{5,\Omega}. \tag{4}$$

We shall construct a nonstandard interpolation operator which improve the optimal estimate (4). Two principal steps are performed for this purpose. The first one is to prove a local supercloseness in energy norm by means of special interpolation operator i_h and the second one is to prove a global ultraconvergence to $I_{2h}P_h u$, where the operator I_{2h} verifies certain ultraconvergence properties.

3. Construction of the Operator $\mathbf{i_h}$

Consider the partition $\tau_h = \bigcup T$, where for any finite element T its center is denoted by (x_T, y_T) and its element dimensions in x and y-direction are denoted by $h_{1,T}, h_{2,T}$, respectively. The following denotations are also adopted:

$$h_T = \max(h_{1,T}; h_{2,T}),\ h = \max_{T \in \tau_h} h_T.$$

Also, the vertices and edges of any element T are noted by s_j and $l_j,\ j = 1, \ldots, 4$, respectively.

Taking in consideration the case $n = 4$ we introduce the interpolation operator $i_h : C^1(T) \to V_h(T)$ using the so-called "vertices - edges" conditions: $\forall v \in C^1(T)$ and $j = 1, \ldots, 4$

$$i_h v(s_j) = v(s_j),\ \frac{\partial i_h v}{\partial x}(s_j) = \frac{\partial v}{\partial x}(s_j),$$

$$\frac{\partial i_h v}{\partial y}(s_j) = \frac{\partial v}{\partial y}(s_j),\ \frac{\partial^2 i_h v}{\partial x \partial y}(s_j) = \frac{\partial^2 v}{\partial x \partial y}(s_j),$$

(5)

$$\int_{l_j} i_h v(x,y)\, dl_j = \int_{l_j} v(x,y)\, dl_j,\ \int_{l_j} \frac{\partial i_h v}{\partial \nu}(x,y)\, dl_j = \int_{l_j} \frac{\partial v}{\partial \nu}(x,y)\, dl_j,$$

$$\int_T i_h v(x,y)\, dx\, dy = \int_T v(x,y)\, dx\, dy.$$

It is evident that $i_h v \equiv v$ for any $v \in V_h$. Let us remark that

$$\|i_h v - v\|_{2,\Omega} \le C h^3 \|v\|_{5,\Omega}.$$

In one-dimensional case (see [3]) the basis determining the operator i_h is $\{\varphi_i(t)\}_{i=1}^5,\ t \in [-1, 1]$, where:

$$\begin{pmatrix} \varphi_1(t) \\ \varphi_2(t) \\ \varphi_3(t) \\ \varphi_4(t) \\ \varphi_5(t) \end{pmatrix} = \frac{1}{16} \begin{pmatrix} -15 & 4 & 30 & -12 & -7 \\ -5 & 4 & 6 & -4 & -1 \\ 15 & 0 & -30 & 0 & 15 \\ -15 & -4 & 30 & 12 & -7 \\ 5 & 4 & -6 & -4 & 1 \end{pmatrix} \begin{pmatrix} t^4 \\ t^3 \\ t^2 \\ t \\ 1 \end{pmatrix}.$$

Then, for the two-dimensional case the basis is obtained by the products $\varphi_i(x).\varphi_s(y),\ i, s = 1, \ldots, 5$, i.e. is $\{\psi_j(x,y)\}_{j=1}^{25}$. Moreover,

$$i_h v(x,y)|_T = \sum_{j=1}^{25} \hat{v}_j . \psi_j \left(\frac{x - x_T}{h_{1,T}}, \frac{y - y_T}{h_{2,T}} \right),$$

where

$$\hat{v}_{4i-3} = v(s_i),\ \hat{v}_{4i-2} = h_{1,T} \frac{\partial v}{\partial x}(s_i),$$

$$\hat{v}_{4i-1} = h_{2,T} \frac{\partial v}{\partial y}(s_i),\ \hat{v}_{4i} = h_{1,T} h_{2,T} \frac{\partial^2 v}{\partial x \partial y}(s_i),\ i = 1, \ldots, 4;$$

$$\hat{v}_{2i+15} = \frac{1}{h_{1,T}} \int_{l_i} v(x,y)\, dx,\ \hat{v}_{2i+16} = \frac{h_{2,T}}{h_{1,T}} \int_{l_i} \frac{\partial v}{\partial y}(x,y)\, dx,\ i = 1; 3;$$

$$\hat{v}_{2i+15} = \frac{1}{h_{2,T}} \int_{l_i} v(x,y)\, dy,\ \hat{v}_{2i+16} = \frac{h_{1,T}}{h_{2,T}} \int_{l_i} \frac{\partial v}{\partial x}(x,y)\, dy,\ i = 2; 4;$$

$$\hat{v}_{25} = \frac{1}{h_{1,T} h_{2,T}} \int_T v(x,y)\, dx\, dy.$$

Remark 1. *It is easy to see that i_h satisfies the C^1-condition. That is continuity of the function $i_h v$ and continuity of its first derivatives across the interelement boundaries.*

4. Superclose Properties of $\mathbf{i_h}$

This section presents important error estimates which are related to the interpolated finite element method. The difficulties for fourth-order elliptic problems arise from the availability of the mixed derivatives in the variational bilinear $a-$form. However,

$$a(u,v)=\int_\Omega\left[\frac{\partial^2 u}{\partial x^2}\frac{\partial^2 v}{\partial x^2}+2\frac{\partial^2 u}{\partial x\partial y}\frac{\partial^2 v}{\partial x\partial y}+\frac{\partial^2 u}{\partial y^2}\frac{\partial^2 v}{\partial y^2}\right]dx\,dy.$$

This presentation is not unique in terms of subject-matter (see for example [10]).

Let us introduce the following error functions for any $T\in\tau_h$:

$$E(x)=\frac{1}{2}\left[(x-x_T)^2-h_{1,T}^2\right],$$

$$F(y)=\frac{1}{2}\left[(y-y_T)^2-h_{2,T}^2\right].$$

We use the identities:

$$\begin{aligned} x-x_T &= \frac{1}{90}\left[E^3(x)\right]^{(5)},\\ (x-x_T)^2 &= \frac{1}{1260}\left[E^4(x)\right]^{(6)}+\frac{h_{1,T}^2}{7}. \end{aligned}\tag{6}$$

Further, we use another pair of identities:

$$\begin{aligned} (x-x_T)^3 &= \frac{1}{420}\left[E^4(x)\right]^{(5)} + \frac{h_{1,T}^2}{210}\left[E^3(x)\right]^{(5)},\\ (x-x_T)^4 &= \frac{1}{4725}\left[E^5(x)\right]^{(6)} + \frac{h_{1,T}^2}{1890}\left[E^4(x)\right]^{(6)}+\frac{h_{1,T}^4}{21}. \end{aligned}\tag{7}$$

The following Lemma gives some integral presentations by means of error function $E(x)$:

Lemma 1. *Let $v(x,y)$ be a sufficiently smooth function defined on $\overline{T}, T\in\tau_h$. Then*

(i): *If* $\int_{l_i} v(x,y)\,dy=\int_{l_i}\frac{\partial v}{\partial x}(x,y)\,dy=0,\ i=2;4,$ *then*

$$\begin{aligned} \int_T (x-x_T)v(x,y)\,dx\,dy &= -\frac{1}{90}\int_T E^3(x)\frac{\partial^5 v}{\partial x^5}(x,y)\,dx\,dy;\\ \int_T (x-x_T)^3 v(x,y)\,dx\,dy &= -\frac{1}{420}\int_T E^4(x)\frac{\partial^5 v}{\partial x^5}(x,y)\,dx\,dy\\ &\quad -\frac{h_{1,T}^2}{210}\int_T E^3(x)\frac{\partial^5 v}{\partial x^5}(x,y)\,dx\,dy; \end{aligned}\tag{8}$$

(ii): *If* $\int_{l_i} v(x,y)\,dy = \int_{l_i} \frac{\partial v}{\partial x}(x,y)\,dy = \int_T v(x,y)\,dx\,dy = 0,\ i = 2;4,$ *then*

$$\begin{aligned}
\int_T (x-x_T)^2 v(x,y)\,dx\,dy &= \frac{1}{1260}\int_T E^4(x)\frac{\partial^6 v}{\partial x^6}(x,y)\,dx\,dy; \\
\int_T (x-x_T)^4 v(x,y)\,dx\,dy &= \frac{1}{4725}\int_T E^5(x)\frac{\partial^6 v}{\partial x^6}(x,y)\,dx\,dy \\
&\quad + \frac{h_{1,T}^2}{1890}\int_T E^4(x)\frac{\partial^6 v}{\partial x^6}(x,y)\,dx\,dy.
\end{aligned} \tag{9}$$

Proof. The four equalities (8) and (9) are proved by using identities (6) and (7) for the terms $(x-x_T)^i,\ i = 1,\ldots,4$. The Green's formula and the above conditions are applied. □

Next, other identities related to the two error functions $E(x)$ and $F(y)$ are needed:

$$\begin{aligned}
(x-x_T)^3 &= \frac{1}{15}\left[E^3(x)\right]^{(3)} + \frac{3h_{1,T}^2}{5}(x-x_T), \\
(x-x_T)^4 &= \frac{1}{105}\left[E^4(x)\right]^{(4)} + \frac{6h_{1,T}^2}{7}(x-x_T)^2 - \frac{3h_{1,T}^4}{35},
\end{aligned} \tag{10}$$

and correspondingly

$$\begin{aligned}
(y-y_T)^3 &= \frac{1}{15}\left[F^3(y)\right]^{(3)} + \frac{3h_{2,T}^2}{5}(y-y_T), \\
(y-y_T)^4 &= \frac{1}{105}\left[F^4(y)\right]^{(4)} + \frac{6h_{2,T}^2}{7}(y-y_T)^2 - \frac{3h_{2,T}^4}{35}.
\end{aligned} \tag{11}$$

Lemma 2. *Suppose that for sufficiently smooth function* $v(x,y)$

$$\frac{\partial^{i+j} v}{\partial x^i \partial y^j}(s_k) = 0,\ i,j = 0,1;$$

$$\int_{l_k} v\,dl = \int_{l_k} \frac{\partial v}{\partial \nu}\,dl = \int_T v\,dx\,dy = 0,\ k = 1,\ldots,4,$$

then the following six relations for the function $(x-x_T)^i(y-y_T)^j\dfrac{\partial^4 v}{\partial x^2\partial y^2}$ *when* $i,j \in \{0,\ldots,4\}$ *are valid:*

($\mathbf{R_1}$) *For* $i,j = 0,1,2;$

$$\int_T (x-x_T)^i(y-y_T)^j\frac{\partial^4 v}{\partial x^2\partial y^2}\,dx\,dy = 0;$$

($\mathbf{R_2}$) *For $j = 0, 1, 2$ and, by analogy for $i = 0, 1, 2$*

$$\int_T (x - x_T)^3 (y - y_T)^j \frac{\partial^4 v}{\partial x^2 \partial y^2}\, dx\, dy = -\frac{1}{15} \int_T E^3(x)(y - y_T)^j \frac{\partial^7 v}{\partial x^5 \partial y^2}\, dx\, dy,$$
$$\int_T (x - x_T)^i (y - y_T)^3 \frac{\partial^4 v}{\partial x^2 \partial y^2}\, dx\, dy = -\frac{1}{15} \int_T F^3(y)(x - x_T)^i \frac{\partial^7 v}{\partial x^2 \partial y^5}\, dx\, dy;$$

($\mathbf{R_3}$) *For $j = 0, 1, 2$ and $i = 0, 1, 2$, respectively*

$$\int_T (x - x_T)^4 (y - y_T)^j \frac{\partial^4 v}{\partial x^2 \partial y^2}\, dx\, dy = \frac{1}{105} \int_T E^4(x)(y - y_T)^j \frac{\partial^8 v}{\partial x^6 \partial y^2}\, dx\, dy,$$
$$\int_T (x - x_T)^i (y - y_T)^4 \frac{\partial^4 v}{\partial x^2 \partial y^2}\, dx\, dy = \frac{1}{105} \int_T F^4(y)(x - x_T)^i \frac{\partial^8 v}{\partial x^2 \partial y^6}\, dx\, dy;$$

($\mathbf{R_4}$) *For $i, j = 3$*

$$\int_T (x - x_T)^3 (y - y_T)^3 \frac{\partial^4 v}{\partial x^2 \partial y^2}\, dx\, dy = -\frac{1}{15} \int_T E^3(x)(y - y_T)^3 \frac{\partial^7 v}{\partial x^5 \partial y^2}\, dx\, dy$$
$$-\frac{h_{1,T}^2}{25} \int_T F^3(y)(x - x_T) \frac{\partial^7 v}{\partial x^2 \partial y^5}\, dx\, dy;$$

($\mathbf{R_5}$) *For $(i, j) = (3, 4)$ and $(i, j) = (4, 3)$, respectively*

$$\int_T (x - x_T)^4 (y - y_T)^3 \frac{\partial^4 v}{\partial x^2 \partial y^2}\, dx\, dy = \frac{1}{105} \int_T E^4(x)(y - y_T)^3 \frac{\partial^7 v}{\partial x^6 \partial y^2}\, dx\, dy$$
$$-\frac{2h_{1,T}^2}{35} \int_T (x - x_T)^2 F^3(y) \frac{\partial^7 v}{\partial x^2 \partial y^5}\, dx\, dy + \frac{h_{1,T}^4}{175} \int_T F^3(y) \frac{\partial^7 v}{\partial x^2 \partial y^5}\, dx\, dy;$$

$$\int_T (x - x_T)^3 (y - y_T)^4 \frac{\partial^4 v}{\partial x^2 \partial y^2}\, dx\, dy = \frac{1}{105} \int_T F^4(y)(x - x_T)^3 \frac{\partial^7 v}{\partial x^2 \partial y^6}\, dx\, dy$$
$$-\frac{2h_{2,T}^2}{35} \int_T (y - y_T)^2 E^3(x) \frac{\partial^7 v}{\partial x^5 \partial y^2}\, dx\, dy + \frac{h_{2,T}^4}{175} \int_T E^3(x) \frac{\partial^7 v}{\partial x^5 \partial y^2}\, dx\, dy;$$

($\mathbf{R_6}$) *For $i, j = 4$*

$$\int_T (x - x_T)^4 (y - y_T)^4 \frac{\partial^4 v}{\partial x^2 \partial y^2}\, dx\, dy = \frac{1}{105} \int_T E^4(x)(y - y_T)^4 \frac{\partial^7 v}{\partial x^6 \partial y^2}\, dx\, dy$$

$$+\frac{2h_{1,T}^2}{245}\int_T (x-x_T)^2F^4(y)\frac{\partial^7 v}{\partial x^2\partial y^6}\,dx\,dy-\frac{h_{1,T}^4}{1225}\int_T F^4(y)\frac{\partial^8 v}{\partial x^2\partial y^6}\,dx\,dy.$$

Proof. Direct calculations proves the first relation. The next one follows by presentations of $(x-x_T)^3$ and $(y-y_T)^3$ from (10), (11). Then integrating by parts and using (R_1) we obtain (R_2).

By way of analogy with the previous case the relation (R_3) is connected with the presentation of $(x-x_T)^4$ and $(y-y_T)^4$ from (10), (11).

In order to prove the cases (R_4) and (R_5) we transform $(x-x_T)^3$ and $(x-x_T)^4$ by means of (10). Then, after integrating by parts we use the relation (R_2).

The last case (R_6) follows from the presentation of $(x-x_T)^4$ in (10) and from the corresponding equality of (R_3). □

Finally, the last two lemmas make it possible to formulate a superclose property concerning the interpolation operator i_h. This important result will be proved in the following theorem:

Theorem 1. *Let the solution $u(x,y)$ of problem (3) belongs to the space $H^7(\Omega)$. Then, for any $v\in V_h(\Omega)$*

$$a(i_hu-u,v)\le Ch^5\|u\|_{7,\Omega}\|v\|_{2,\Omega}. \tag{12}$$

Proof. Let us denote $U=i_hu-u$. The three ingredients of the $a-$form $a(U,v)$ have to be estimated. It should be noted that both terms

$$\int_\Omega \frac{\partial^2 U}{\partial x^2}\frac{\partial^2 v}{\partial x^2}\,dx\,dy \text{ and } \int_\Omega \frac{\partial^2 U}{\partial y^2}\frac{\partial^2 v}{\partial y^2}\,dx\,dy$$

can be treated in a similar way.

Step 1 Consider $\int_\Omega \frac{\partial^2 U}{\partial y^2}\frac{\partial^2 v}{\partial y^2}\,dx\,dy,\ \forall v\in V_h$. So, for $(x,y)\in T$, the second factor is expanded as follows:

$$\frac{\partial^2 v}{\partial y^2}(x,y)=\sum_{s=0}^{4}\frac{(x-x_T)^s}{s!}\frac{\partial^{s+2}v}{\partial x^s\partial y^2}(x_T,y),$$

taking into account that $v(x,y)|_T\in Q_4(T)$, consequently

$$\int_\Omega \frac{\partial^2 U}{\partial y^2}\frac{\partial^2 v}{\partial y^2}\,dx\,dy=\sum_{T\in\tau_h}\sum_{s=0}^{4}\int_T\frac{(x-x_T)^s}{s!}\frac{\partial^2 U}{\partial y^2}\frac{\partial^{s+2}v}{\partial x^s\partial y^2}(x_T,y)\,dx\,dy. \tag{13}$$

The first term ($s=0$) in the right-hand side of (13) is cancelled. This fact follows by definition of $U(x,y)$ and using Green's formula, i.e.

$$\int_T \frac{\partial^2 U}{\partial y^2}\frac{\partial^2 v}{\partial y^2}(x_T,y)\,dx\,dy=0. \tag{14}$$

In order to transform the other components of (13), it has to be verified that the functions

$$\frac{\partial^2 U}{\partial y^2}\frac{\partial^{s+2} v}{\partial x^s \partial y^2}(x_T, y),\ s = 1, \ldots, 4,$$

fulfill the conditions of Lemma 1. Hence, the relations (8) and (9) will be satisfied for $s = 1; 3$ and $s = 2; 4$ respectively.

Let $i = 2; 4$ and $s = 1, \ldots, 4$. Integrating by parts reveals that:

$$\begin{aligned}
\int_{l_i} \frac{\partial^2 U}{\partial y^2}\frac{\partial^{s+2} v}{\partial x^s \partial y^2}(x_T, y)\, dy &= \frac{\partial U}{\partial y}(x_T \pm h_{1,T}, t)\ \frac{\partial^{s+2} v}{\partial x^s \partial y^2}(x_T, y)\Big|_{y_T - h_{2,T}}^{y_T + h_{2,T}} \\
-\int_{l_i} \frac{\partial U}{\partial y}\frac{\partial^{s+3} v}{\partial x^s \partial y^3}(x_T, y)\, dy &= -U(x_T \pm h_{1,T}, y)\ \frac{\partial^{s+3} v}{\partial x^s \partial y^3}(x_T, y)\Big|_{y_T - h_{2,T}}^{y_T + h_{2,T}} \qquad (15) \\
+\int_{l_i} U\frac{\partial^{s+4} v}{\partial x^s \partial y^4}(x_T, y)\, dy &= 0.
\end{aligned}$$

By analogy, it is easy to get the equality:

$$\int_{l_i} \frac{\partial}{\partial x}\left[\frac{\partial^2 U}{\partial y^2}\frac{\partial^{s+2} v}{\partial x^s \partial y^2}(x_T, y)\right] dy = \int_{l_i} \frac{\partial^3 U}{\partial x \partial y^2}\frac{\partial^{s+2} v}{\partial x^s \partial y^2}(x_T, y)\, dy = 0, \qquad (16)$$

having in mind the definition of U.

For $s = 2; 4$ and using the Green's formula we get:

$$\begin{aligned}
\int_T \frac{\partial^2 U}{\partial y^2}\frac{\partial^{s+2} v}{\partial x^s \partial y^2}(x_T, y)\, dx\, dy &= \left(\int_{l_3} - \int_{l_1}\right) \frac{\partial U}{\partial y}\frac{\partial^{s+2} v}{\partial x^s \partial y^2}(x_T, y)\, dx \\
-\int_T \frac{\partial U}{\partial y}\frac{\partial^{s+3} v}{\partial x^s \partial y^3}(x_T, y)\, dx\, dy &= \left(\int_{l_3} - \int_{l_1}\right) U\frac{\partial^{s+3} v}{\partial x^s \partial y^3}(x_T, y)\, dx \qquad (17) \\
+\int_T U\frac{\partial^{s+4} v}{\partial x^s \partial y^4}(x_T, y)\, dx\, dy &= 0.
\end{aligned}$$

Then, in view of (15) and (16), applying (8) of Lemma 1 for the function $\dfrac{\partial^2 U}{\partial y^2}\dfrac{\partial^{s+2} v}{\partial x^s \partial y^2}(x_T, y)$ with $s = 1$ and $s = 3$ respectively, we obtain:

$$\begin{aligned}
&\int_T (x - x_T)\frac{\partial^2 U}{\partial y^2}\frac{\partial^3 v}{\partial x \partial y^2}(x_T, y)\, dx\, dy \\
&= \frac{1}{90}\int_T E^3(x)\frac{\partial^7 u}{\partial x^5 \partial y^2}\frac{\partial^3 v}{\partial x \partial y^2}(x_T, y)\, dx\, dy;
\end{aligned} \qquad (18)$$

$$\int_T \frac{(x-x_T)^3}{3!}\frac{\partial^2 U}{\partial y^2}\frac{\partial^5 v}{\partial x^3\partial y^2}(x_T,y)\,dx\,dy$$

$$= \frac{1}{420.3!}\int_T E^4(x)\frac{\partial^7 u}{\partial x^5\partial y^2}\frac{\partial^5 v}{\partial x^3\partial y^2}(x_T,y)\,dx\,dy \tag{19}$$

$$+\frac{h_{1,T}^2}{210.3!}\int_T E^3(x)\frac{\partial^7 u}{\partial x^5\partial y^2}\frac{\partial^5 v}{\partial x^3\partial y^2}(x_T,y)\,dx\,dy.$$

Similarly, we get:

$$\int_T \frac{(x-x_T)^2}{2!}\frac{\partial^2 U}{\partial y^2}\frac{\partial^4 v}{\partial x^2\partial y^2}(x_T,y)\,dx\,dy$$

$$= -\frac{1}{1260.2!}\int_T E^4(x)\frac{\partial^8 u}{\partial x^6\partial y^2}\frac{\partial^4 v}{\partial x^2\partial y^2}(x_T,y)\,dx\,dy;$$

$$\int_T \frac{(x-x_T)^4}{4!}\frac{\partial^2 U}{\partial y^2}\frac{\partial^6 v}{\partial x^4\partial y^2}(x,y)\,dx\,dy$$

$$= -\frac{1}{4725.4!}\int_T E^5(x)\frac{\partial^8 u}{\partial x^6\partial y^2}\frac{\partial^6 v}{\partial x^4\partial y^2}(x,y)\,dx\,dy$$

$$-\frac{h_{1,T}^2}{1890.4!}\int_T E^4(x)\frac{\partial^8 u}{\partial x^6\partial y^2}\frac{\partial^6 v}{\partial x^4\partial y^2}(x,y)\,dx\,dy,$$

having in mind (15)-(17) and applying (9) of Lemma 1 for the same function with $s=2$ and $s=4$ respectively.

It is possible to reduce the requirement for high smoothness without loss of accuracy. Indeed, using the identity

$$\left[E^k(x)\right]' = kE^{k-1}(x).(x-x_T),\ k=4;5,$$

the last two equalities may be written in the form

$$\int_T \frac{(x-x_T)^2}{2!}\frac{\partial^2 U}{\partial y^2}\frac{\partial^4 v}{\partial x^2\partial y^2}(x_T,y)\,dx\,dy$$

$$= \frac{1}{315.2!}\int_T E^3(x)(x-x_T)\frac{\partial^7 u}{\partial x^5\partial y^2}\frac{\partial^4 v}{\partial x^2\partial y^2}(x_T,y)\,dx\,dy; \tag{20}$$

$$
\begin{aligned}
&\int_T \frac{(x-x_T)^4}{4!}\frac{\partial^2 U}{\partial y^2}\frac{\partial^6 v}{\partial x^4\partial y^2}(x,y)\,dx\,dy \\
&= \frac{1}{945.4!}\int_T E^4(x)(x-x_T)\frac{\partial^7 u}{\partial x^5\partial y^2}\frac{\partial^6 v}{\partial x^4\partial y^2}(x,y)\,dx\,dy \qquad (21)\\
&+\frac{h_{1,T}^2}{1890.3!}\int_T E^3(x)(x-x_T)\frac{\partial^7 u}{\partial x^5\partial y^2}\frac{\partial^6 v}{\partial x^4\partial y^2}(x,y)\,dx\,dy.
\end{aligned}
$$

The right-hand side of (18), (19) and (20) can be transformed for $v \in \tau_h$ by the identity

$$\frac{\partial^{s+2} v}{\partial x^s \partial y^2}(x_T,y) = \sum_{i=s}^{4}\frac{(x_T-x)^{i-s}}{(i-s)!}\frac{\partial^{i+2} v}{\partial x^i\partial y^2}(x,y),\ s=1,2,3;\ (x,y)\in T.$$

From (14), (18)-(21), using the inverse inequality [5], we obtain

$$\int_T \frac{\partial^2 U}{\partial y^2}\frac{\partial^2 v}{\partial y^2}\,dx\,dy \le Ch^5\|u\|_{7,T}\|v\|_{2,T},$$

consequently,

$$\int_\Omega \frac{\partial^2 U}{\partial y^2}\frac{\partial^2 v}{\partial y^2}\,dx\,dy \le Ch^5\|u\|_{7,\Omega}\|v\|_{2,\Omega}, \qquad (22)$$

and by analogy

$$\int_\Omega \frac{\partial^2 U}{\partial x^2}\frac{\partial^2 v}{\partial x^2}\,dx\,dy \le Ch^5\|u\|_{7,\Omega}\|v\|_{2,\Omega}. \qquad (23)$$

Step 2 Here we consider the mixed term of the $a-$form. Let's consider that the following equalities hold:

$$
\begin{aligned}
\sum_{T\in\tau_h}\int_T \frac{\partial^2 U}{\partial x\partial y}\frac{\partial^2 v}{\partial x\partial y}\,dx\,dy &= \sum_{T\in\tau_h}\left(\int_{l_3}-\int_{l_1}\right)\frac{\partial^2 U}{\partial x\partial y}\frac{\partial v}{\partial x}\,dx \\
-\sum_{T\in\tau_h}\int_T \frac{\partial^3 U}{\partial x\partial y^2}\frac{\partial v}{\partial x}\,dx\,dy &= -\sum_{T\in\tau_h}\left(\int_{l_2}-\int_{l_4}\right)\frac{\partial^3 U}{\partial x\partial y^2}v\,dx \qquad (24)\\
+\sum_{T\in\tau_h}\int_T \frac{\partial^4 U}{\partial x^2\partial y^2}v\,dx\,dy &= \sum_{T\in\tau_h}\int_T \frac{\partial^4 U}{\partial x^2\partial y^2}v\,dx\,dy,
\end{aligned}
$$

because the functions $\dfrac{\partial^2 U}{\partial x\partial y}\dfrac{\partial v}{\partial y}$ and $\dfrac{\partial^3 U}{\partial x\partial y^2}v$ are continuous on the edges, parallel to the x and $y-$axes, respectively. For the line integrals we integrate over the common side of each pair of two adjacent elements with the same integrands, but in opposite direction. For integrals over any parts of Γ the function v vanishes.

Consider the following presentation for any $v \in V_h$:

$$v(x,y) = \sum_{i,j=0}^{4} \frac{1}{i!j!}(x-x_T)^i(y-y_T)^j \frac{\partial^{i+j} v}{\partial x^i \partial y^j}(x_T, y_T) \ \forall (x,y) \in T.$$

Combining this equality with (24) we get:

$$\int_T \frac{\partial^4 U}{\partial x^2 \partial y^2} v \, dx \, dy = \sum_{i,j=0}^{4} \int_T \frac{(x-x_T)^i(y-y_T)^j}{i!j!} \frac{\partial^4 U}{\partial x^2 \partial y^2} \frac{\partial^{i+j} v}{\partial x^i \partial y^j}(x_T, y_T) \, dx \, dy. \tag{25}$$

We shall estimate each term in (25) using the relations of Lemma 2.
First, from (R_1) for $i, j = 0, 1, 2$ it follows that

$$\int_T \frac{(x-x_T)^i(y-y_T)^j}{i!j!} \frac{\partial^4 U}{\partial x^2 \partial y^2} \frac{\partial^{i+j} v}{\partial x^i \partial y^j}(x_T, y_T) \, dx \, dy = 0. \tag{26}$$

Next, the relation (R_2) for $i; j = 0, 1, 2$ respectively gives

$$\begin{aligned} &\int_T \frac{(x-x_T)^3(y-y_T)^j}{3!j!} \frac{\partial^4 U}{\partial x^2 \partial y^2} \frac{\partial^{3+j} v}{\partial x^3 \partial y^j}(x_T, y_T) \, dx \, dy \\ &= \frac{1}{15.3!j!} \int_T E^3(x)(y-y_T)^j \frac{\partial^7 u}{\partial x^5 \partial y^2} \frac{\partial^{3+j} v}{\partial x^3 \partial y^j}(x_T, y_T) \, dx \, dy; \end{aligned}$$

$$\begin{aligned} &\int_T \frac{(x-x_T)^i(y-y_T)^3}{i!3!} \frac{\partial^4 U}{\partial x^2 \partial y^2} \frac{\partial^{i+3} v}{\partial x^i \partial y^3}(x_T, y_T) \, dx \, dy \\ &= \frac{1}{15.i!3!} \int_T (x-x_T)^i F^3(y) \frac{\partial^7 u}{\partial x^2 \partial y^5} \frac{\partial^{i+3} v}{\partial x^i \partial y^3}(x_T, y_T) \, dx \, dy. \end{aligned} \tag{27}$$

By analogy with (27), using (R_3) for $i; j = 0, 1, 2$, we obtain the next two equalities:

$$\begin{aligned} &\int_T \frac{(x-x_T)^4(y-y_T)^j}{4!j!} \frac{\partial^4 U}{\partial x^2 \partial y^2} \frac{\partial^{4+j} v}{\partial x^4 \partial y^j}(x_T, y_T) \, dx \, dy \\ &= -\frac{1}{105.4!j!} \int_T E^4(x)(y-y_T)^j \frac{\partial^8 u}{\partial x^6 \partial y^2} \frac{\partial^{4+j} v}{\partial x^4 \partial y^j}(x_T, y_T) \, dx \, dy \\ &= \frac{1}{105.3!j!} \int_T E^3(x)(x-x_T)(y-y_T)^j \frac{\partial^7 u}{\partial x^5 \partial y^2} \frac{\partial^{4+j} v}{\partial x^4 \partial y^j}(x_T, y_T) \, dx \, dy; \end{aligned}$$

$$\begin{aligned} &\int_T \frac{(x-x_T)^i(y-y_T)^4}{i!4!} \frac{\partial^4 U}{\partial x^2 \partial y^2} \frac{\partial^{i+4} v}{\partial x^i \partial y^4}(x_T, y_T) \, dx \, dy \\ &= -\frac{1}{105.i!4!} \int_T (x-x_T)^i F^4(y) \frac{\partial^8 u}{\partial x^2 \partial y^6} \frac{\partial^{i+4} v}{\partial x^i \partial y^4}(x_T, y_T) \, dx \, dy \\ &= \frac{1}{105.i!3!} \int_T (x-x_T)^i (y-y_T) F^3(y) \frac{\partial^7 u}{\partial x^2 \partial y^5} \frac{\partial^{i+4} v}{\partial x^i \partial y^4}(x_T, y_T) \, dx \, dy. \end{aligned} \tag{28}$$

From (R_4) it follows that

$$\int_T \frac{(x-x_T)^3(y-y_T)^3}{3!3!}\frac{\partial^4 U}{\partial x^2\partial y^2}\frac{\partial^6 v}{\partial x^3\partial y^3}(x_T,y_T)\,dx\,dy$$

$$= \frac{1}{15.3!.3!}\int_T E^3(x)(y-y_T)^3\frac{\partial^7 u}{\partial x^5\partial y^2}\frac{\partial^6 v}{\partial x^3\partial y^3}(x_T,y_T)\,dx\,dy. \tag{29}$$

$$+\frac{h_{1,T}^2}{25.3!3!}\int_T (x-x_T)F^3(y)\frac{\partial^7 u}{\partial x^2\partial y^5}\frac{\partial^6 v}{\partial x^3\partial y^3}(x_T,y_T)\,dx\,dy.$$

Introducing the equalities for sufficiently smooth function U

$$\int_T E^4(x)\frac{\partial^6 U}{\partial x^6}\,dx\,dy = -4\int_T (x-x_T)E^3(x)\frac{\partial^5 U}{\partial x^5}\,dx\,dy,$$

$$\int_T F^4(y)\frac{\partial^6 U}{\partial y^6}\,dx\,dy = -4\int_T (y-y_T)F^3(y)\frac{\partial^5 U}{\partial y^5}\,dx\,dy,$$

from the relation (R_4) we get

$$\int_T \frac{(x-x_T)^4(y-y_T)^3}{4!3!}\frac{\partial^4 U}{\partial x^2\partial y^2}\frac{\partial^7 v}{\partial x^4\partial y^3}(x_T,y_T)\,dx\,dy$$

$$= \frac{1}{105.3!.3!}\int_T E^3(x)(x-x_T)(y-y_T)^3\frac{\partial^7 u}{\partial x^5\partial y^2}\frac{\partial^7 v}{\partial x^4\partial y^3}(x_T,y_T)\,dx\,dy. \tag{30}$$

$$+\frac{2h_{1,T}^2}{35.4!3!}\int_T (x-x_T)^2F^3(y)\frac{\partial^7 u}{\partial x^2\partial y^5}\frac{\partial^7 v}{\partial x^4\partial y^3}(x_T,y_T)\,dx\,dy$$

$$-\frac{h_{1,T}^4}{175.4!3!}\int_T F^3(y)\frac{\partial^7 u}{\partial x^2\partial y^5}\frac{\partial^7 v}{\partial x^4\partial y^3}(x_T,y_T)\,dx\,dy.$$

Similarly,

$$\int_T \frac{(x-x_T)^3(y-y_T)^4}{3!4!}\frac{\partial^4 U}{\partial x^2\partial y^2}\frac{\partial^7 v}{\partial x^3\partial y^4}(x_T,y_T)\,dx\,dy$$

$$= \frac{1}{105.3!.3!}\int_T (x-x_T)^3(y-y_T)F^3(y)\frac{\partial^7 u}{\partial x^2\partial y^5}\frac{\partial^7 v}{\partial x^3\partial y^4}(x_T,y_T)\,dx\,dy. \tag{31}$$

$$+\frac{2h_{2,T}^2}{35.3!4!}\int_T E^3(x)(y-y_T)^2\frac{\partial^7 u}{\partial x^5\partial y^2}\frac{\partial^7 v}{\partial x^3\partial y^4}(x_T,y_T)\,dx\,dy$$

$$-\frac{h_{2,T}^4}{175.3!4!}\int_T E^3(x)\frac{\partial^7 u}{\partial x^5\partial y^2}\frac{\partial^7 v}{\partial x^3\partial y^4}(x_T,y_T)\,dx\,dy.$$

By analogy

$$\int_T \frac{(x-x_T)^4(y-y_T)^4}{4!4!}\frac{\partial^4 U}{\partial x^2\partial y^2}\frac{\partial^8 v}{\partial x^4\partial y^4}\,dx\,dy$$

$$= \frac{1}{105.4!.3!}\int_T E^3(x)(x-x_T)(y-y_T)^4\frac{\partial^7 u}{\partial x^5\partial y^2}\frac{\partial^8 v}{\partial x^4\partial y^4}\,dx\,dy. \tag{32}$$

$$+\frac{2h_{1,T}^2}{245.4!4!}\int_T (x-x_T)^2(y-y_T)F^3(y)\frac{\partial^7 u}{\partial x^2\partial y^5}\frac{\partial^8 v}{\partial x^4\partial y^4}\,dx\,dy$$
$$-\frac{h_{1,T}^4}{1225.4!4!}\int_T (y-y_T)F^3(y)\frac{\partial^7 u}{\partial x^2\partial y^5}\frac{\partial^8 v}{\partial x^4\partial y^4}\,dx\,dy.$$

In the right-hand sides of (27)-(31) we apply the equality

$$\frac{\partial^{i+j}v}{\partial x^i\partial y^j}(x_T,y_T)=\sum_{i_0=i,j_0=j}^{4}\frac{(x-x_T)^{i_0-i}(y-y_T)^{j_0-j}}{(i_0-i)!(j_0-j)!}\frac{\partial^{i_0+j_0}v}{\partial x^{i_0}\partial y^{j_0}}(x,y),$$

for corresponding values of i and j.

Thus, from (26)-(32) and using the inverse inequality, we obtain

$$\int_T \frac{\partial^2 U}{\partial x\partial y}\frac{\partial^2 v}{\partial x\partial y}\,dx\,dy\le Ch_T^5\|u\|_{7,T}\|v\|_{2,T},$$

and consequently,

$$\int_\Omega \frac{\partial^2 U}{\partial x\partial y}\frac{\partial^2 v}{\partial x\partial y}\,dx\,dy\le Ch^5\|u\|_{7,\Omega}\|v\|_{2,\Omega}. \tag{33}$$

Accordingly, the three terms of $a-$form are estimated. Hence (22), (23) and (33) prove the theorem. □

As a consequence of this theorem the following ultraconvergence result holds:

Theorem 2. *Let the conditions of Theorem 1 be fulfilled. Then*

$$\|i_h u-P_h u\|_{2,\Omega}\le Ch^5\|u\|_{7,\Omega}. \tag{34}$$

Proof. It follows that ($\alpha=const$):

$$\begin{aligned}
\alpha\|i_h u-P_h u\|_{2,\Omega}^2 &\le a(i_h u-P_h u,i_h u-P_h u) && (V\text{-ellipticity})\\
&= a(i_h u-u,i_h u-P_h u) && (\text{elliptic projector})\\
&\le C\|i_h u-u\|_{2,\Omega}\|i_h u-P_h u\|_{2,\Omega} && (\text{continuity})\\
&\le Ch^5\|u\|_{7,\Omega}\|i_h u-P_h u\|_{2,\Omega} && (\text{from (33)}).
\end{aligned}$$

□

5. High Interpolation Operator $\mathbf{I_{2h}}$ - Construction and Estimations

First, let us consider the one-dimensional case. We keep the same notation for one-dimensional FE partition, namely $\tau_h=\{[p_{i-1},p_i],\ i=1,\dots,n\}$. Without loss of generality one can suppose that n is an even natural number. Then the elements of τ_h are combined by pairs of adjacent elements. A new FE partition is defined by

$$\widetilde{\tau}_{2h}=\left\{[p_{2k-2},p_{2k-1}]\cup[p_{2k-1},p_{2k}],\ k=1,\dots,\frac{n}{2}\right\}.$$

The interpolation operator I_{2h} is characterized by the following conditions for every finite element belonging to $\widetilde{\tau}_{2h}$ and every smooth function v:

$$I_{2h}v(p_j) = v(p_j),\ j = 2k-2, 2k-1, 2k;$$
$$[I_{2h}v]'(p_j) = v'(p_j),\ j = 2k-2, 2k-1, 2k;$$
$$\int_{p_j}^{p_{j+1}} I_{2h}v(x)\,dx = \int_{p_j}^{p_{j+1}} v(x)\,dx,\ j = 2k-2, 2k-1.$$

For instance, it is possible to obtain the following basic functions defined on a reference element, i.e. for $t \in [-1,0]\cup[0,1]$:

$$\begin{aligned}
\psi_{1;7}(t) &= \frac{1}{16}t^2(1-t)^2(\pm 72t^3 + 95t^2 \mp 14t - 33),\\
\psi_{2;8}(t) &= \frac{1}{16}t^2(1-t)^2(8t^3 \pm 9t^2 - 2t \mp 33),\\
\psi_{3;6}(t) &= \frac{1}{16}t^2(1-t)^2(1+t)^2(35 \mp 64t),\\
\psi_4(t) &= (1-t)^2(1+t)^2(1-7t^2),\\
\psi_5(t) &= t(1-t)^2(1+t)^2(1-2t)(1+2t).
\end{aligned}$$

By affine transformation

$$t = \frac{x - p_{2k-1}}{h},$$

where

$$h = \begin{cases} p_{2k} - p_{2k-1}, & x \in [p_{2k-1}, p_{2k}],\\ p_{2k-1} - p_{2k-2}, & x \in [p_{2k-2}, p_{2k-1}], \end{cases}$$

it is not difficult to obtain the basic functions $\psi_j(x),\ j = 1,\dots,8$ for any element $[p_{2k-2}, p_{2k-1}]\cup[p_{2k-1}, p_{2k}] \in \widetilde{\tau}_{2h}$.

The set of degrees of freedom is $v(p_j),\ hv'(p_j),\ j = 2k-2, 2k-1, 2k$ and $\frac{1}{h}\int_{p_j}^{p_{j+1}} v(x)\,dx,\ j = 2k-2, 2k-1$.

For 2−dimensional case the basic functions are defined as a product of two one-dimensional basic functions, i.e. $\psi_{i,j}(x,y) = \psi_i(x)\psi_j(y),\ i,j = 1,\dots,8$. Consequently, the degrees of freedom are: (i) the values of $v, \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y}$ and $\frac{\partial^2 v}{\partial x \partial y}$ on the vertices of the four subrectangles $T_s \in \tau_h,\ s = 1,\dots,4$ such that $\bigcup_{s=1}^4 T_s \in \widetilde{\tau}_{2h}$; (ii) the integral values of v and $\frac{\partial v}{\partial \nu}$ on the edges l_j of $T_s \in \tau_h,\ s = 1,\dots,4$; (iii) the integral values of v on the four subrectangles $T_s,\ s = 1,\dots,4$ multiplied or devided by the corresponding sizes characterizing the mesh of τ_h.

In this way the operator I_{2h} is defined on the mesh $\widetilde{\tau}_{2h}$ of size $2h$, obtained as a result of arranging in groups of adjacent elements $T_s \in \tau_h,\ s = 1,\dots,4$. Let $\widetilde{V}_{2h} \subset V$ be finite element spaces associated with $\widetilde{\tau}_{2h}$. Then $\widetilde{V}_{2h}$ consists of polynomials of degree 7 at most in each variable.

By constructing I_{2h}, the following properties are valid:

$$I_{2h} \circ i_h = I_{2h}, \tag{35}$$

$$\|I_{2h}v\|_{r,\Omega} \leq C\|v\|_{r,\Omega}\ \forall v \in V_h,\ r = 0, 1, 2, \tag{36}$$

because the interpolation operator $I_{2h} :\ V_h \to \widetilde{V}_{2h}$ is bounded.

Finally, having in mind that the interpolation polynomial $I_{2h}v$ is of degree seven, it follows

$$\|I_{2h}v - v\|_{2,\Omega} \leq Ch^6\|v\|_{7,\Omega}. \tag{37}$$

The next theorem contains the main estimation:

Theorem 3. *Let $u \in H^7(\Omega)$. The following estimate holds:*

$$\|I_{2h} \circ P_h u - u\|_{2,\Omega} \leq Ch^5\|u\|_{7,\Omega}. \tag{38}$$

Proof. Applying (35), brings to

$$I_{2h} \circ P_h u - u = I_{2h} \circ (P_h u - i_h u) + (I_{2h}u - u)\,.$$

From (36) we have

$$\|I_{2h} \circ P_h u - u\|_{2,\Omega} \leq \|P_h u - i_h u\|_{2,\Omega} + \|I_{2h}u - u\|_{2,\Omega}.$$

Using (37) and Theorem 2, we complete the proof. □

6. Ultraconvergence for Biharmonic Eigenvalue Problem

Here we present a direct application of the result obtained in the previous section. Consider the following fourth-order elliptic eigenvalue problem: find $\lambda \in \mathbf{R},\ u(x, y) \in V, u \neq 0$, such that

$$a(u, v) = \lambda(u, v),\ \forall v \in V. \tag{39}$$

Introduce an approximate eigenvalue problem, which corresponds to (39): find $\lambda_h \in \mathbf{R},\ u_h(x, y) \in V_h, u_h \neq 0$, such that

$$a(u_h, v) = \lambda_h(u_h, v),\ \forall v \in V_h. \tag{40}$$

The assumption here is that the FE space V_h uses polynomials of degree four. A more general case is discussed in [2]. That being the case, it is well known (see [4, 8]) that the rate of convergence of FE approximation to the eigenvalues and eigenfunctions is given by the following estimates:

$$|\lambda - \lambda_h| \leq C(\lambda)h^6\|u\|^2_{5,\Omega}, \tag{41}$$

$$\|u - u_h\|_{2,\Omega} \leq C(\lambda)h^3\|u\|_{5,\Omega}. \tag{42}$$

The solutions of (39) and (40) are related to the Rayleigh quotient

$$\lambda = \frac{a(u, u)}{(u, u)} \text{ and } \lambda_h = \frac{a(u_h, u_h)}{(u_h, u_h)}.$$